

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

RECEIVED
MAY 15 2003
TECH CENTER 1600/2300

Application of

Audrey Goddard et al.

Serial No.: 09/964,994

Filed: September 26, 2001

For: **NOVEL POLYPEPTIDES HAVING
SEQUENCE SIMILARITY TO
CYTOKINE RECEPTORS AND
NUCLEIC ACIDS ENCODING
THE SAME**

Group Art Unit: 1646

Examiner: KEMMERER, ELIZABETH

CERTIFICATE OF MAILING

I hereby certify that this correspondence is being deposited with the United States Postal Service with sufficient postage as first class mail in an envelope addressed to: Commissioner of Patents, P.O. Box 1450, Alexandria, Virginia 22313-1450 on

May 7, 2003

Patricia Tobin

AMENDMENT UNDER 37 C.F.R. §1.111

Commissioner of Patents
P.O. Box 1450
Alexandria, Virginia
22313-1450

Sir:

This is in response to the Office Action (Paper No. 12) mailed herein on January 17, 2003, a response to which is due on April 17, 2003. Enclosed herewith is a Petition for an Extension of Time under 37 C.F.R. §1.136(a) for a response for a period of one (1) month from April 17, 2003, up to and including May 17, 2003, and the fee therefor, thereby making this a timely response.

The Commissioner is authorized to charge any fees which may be required, including extension fees, or credit any overpayment to Deposit Account No. 07-0630.

In the Claims:

Please cancel Claims 1-52 without prejudice or disclaimer.

Please add new claims 53-69.

-53. (New) An isolated nucleic acid molecule having at least 80% sequence identity to:

- (a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;
- (b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide;
- (c) the nucleic acid sequence shown as SEQ ID NO:1;
- (d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or
- (e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532; wherein said encoded polypeptide is a receptor for and binds to the ligand polypeptide shown as SEQ ID NO:4.

54. (New) An isolated nucleic acid molecule having at least 85% sequence identity to:

- (a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;
- (b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide;
- (c) the nucleic acid sequence shown as SEQ ID NO:1;
- (d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or
- (e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532; wherein said encoded polypeptide is a receptor for and binds to the ligand polypeptide shown as SEQ ID NO:4.

55. (New) An isolated nucleic acid molecule having at least 90% sequence identity to:

- (a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;
- (b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2,

- lacking its associated signal peptide;
- (c) the nucleic acid sequence shown as SEQ ID NO:1;
 - (d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or
 - (e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532; wherein said encoded polypeptide is a receptor for and binds to the ligand polypeptide shown as SEQ ID NO:4.
56. (New) An isolated nucleic acid molecule having at least 95% sequence identity to:
- (a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;
 - (b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide;
 - (c) the nucleic acid sequence shown as SEQ ID NO:1;
 - (d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or
 - (e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532; wherein said encoded polypeptide is a receptor for and binds to the ligand polypeptide shown as SEQ ID NO:4.
57. (New) An isolated nucleic acid molecule having at least 99% sequence identity to:
- (a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;
 - (b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide;
 - (c) the nucleic acid sequence shown as SEQ ID NO:1;
 - (d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or
 - (e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532; wherein said encoded polypeptide is a receptor for and binds to the ligand polypeptide shown as SEQ ID NO:4.

58. (New) An isolated nucleic acid molecule comprising:

- (a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;
- (b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide;
- (c) the nucleic acid sequence shown as SEQ ID NO:1;
- (d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or
- (e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532.

59. (New) The isolated nucleic acid molecule of Claim 58 comprising a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2.

60. (New) The isolated nucleic acid molecule of Claim 58 comprising a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide.

61. (New) The isolated nucleic acid molecule of Claim 58 comprising the nucleic acid sequence shown as SEQ ID NO:1.

62. (New) The isolated nucleic acid molecule of Claim 58 comprising the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1.

63. (New) The isolated nucleic acid molecule of Claim 58 comprising the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532.

64. (New) An isolated nucleic acid molecule that hybridizes to:

- (a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;
- (b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2,

lacking its associated signal peptide;

- (c) the nucleic acid sequence shown as SEQ ID NO:1;
- (d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or
- (e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532.

65. (New) The isolated nucleic acid molecule of Claim 64, wherein said hybridization occurs under stringent conditions of 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC (sodium chloride/sodium citrate) and 50% formamide at 55°C, followed by a high-stringency wash consisting of 0.1 x SSC containing EDTA at 55°C.

66. (New) A vector comprising the nucleic acid molecule of Claim 1.

67. (New) The vector of Claim 66, wherein said nucleic acid is operably linked to control sequences recognized by a host cell transformed with the vector.

68. (New) A host cell comprising the vector of Claim 66.

69. (New) The host cell of Claim 68, wherein said cell is a CHO cell, an *E. coli* or a yeast cell.--

REMARKS

Claims 1-52 remain pending in the application for consideration. Claims 1-17 are under consideration subject to Applicants' restriction requirement election. Applicants have herein cancelled Claims 1-52 and added new Claims 53-69 so as to more clearly define the subject matter claimed therein. In that the newly added claims do not introduce new subject matter and are supported in the specification as originally filed, their entry is respectfully requested. Specification support for the newly added Claims 53-69 can be found at least as follows: page 2, lines 14-22, 25-27; page 3, lines 1-37; page 4, lines 1-12, 24-36; page 6, lines 31-37; page 7, lines 1-35; page 8, lines 8-20; page 9, lines 26-30; page 18, lines 1-4; page 52, lines 28-37; and EXAMPLES 1, 11, 12 and 13; and the claims as originally filed.

Information Disclosure Statement

Applicant has been invited to submit a new PTO-1449 form listing the foreign patent documents, with dates listed, for consideration. A new PTO-1449 form is herein submitted listing the missing dates which were previously inadvertently omitted.

Objections and Rejections under 35 U.S.C. §112 Second Paragraph

Claims 4, 9, 10, 11, 12, 16 and 17 stand rejected under 35 U.S.C. §112, second paragraph, as allegedly being indefinite for failing to particularly point out and distinctly claim the subject matter which Applicants regard as the invention. Without necessarily agreeing with the propriety of the rejection, Claims 1-52 have been cancelled and new Claims 53-69 have been added so as to more clearly define the subject matter. The newly added claims correct typographical errors in originally filed Claims 4, 9, 10 and 16. In light of newly added Claims 53-69, Applicants respectfully submit that the indefinite rejection of original Claim 17 has been overcome. In addition, the Examiner has objected to the recitation of "stringent conditions" in Claims 11 and 12, without reciting specific conditions in the original claims. Accordingly, newly added Claims 64-65 have been amended to indicate the specific stringent hybridization conditions. In light of the amendment to the claims and the discussion above, Applicants respectfully request reconsideration and withdrawal of the outstanding

rejections under 35 U.S.C. §112 second paragraph.

Objections and Rejections under 35 U.S.C. §101

Claims 1-17 stand rejected under 35 U.S.C. §101 for allegedly lacking a credible, specific and substantial asserted utility or a well established utility. In light of the claim amendments made herein and for the reasons set forth below, Applicants respectfully traverse.

Utility – Legal Standard

According to the Utility Examination Guidelines ("Utility Guidelines"), 66 Fed. Reg. 1092 (2001) an invention complies with the utility requirement of 35 U.S.C. § 101, if it has at least one asserted "specific, substantial, and credible utility" or a "well-established utility." Under the Utility Guidelines, a utility is "specific" when it is particular to the subject matter claimed. For example, it is generally not enough to state that a polypeptide is useful as a diagnostic without also identifying the condition that is to be diagnosed. The requirement of "substantial utility" defines a "real world" use, and derives from the Supreme Court's holding in *Brenner v. Manson*, 383 U.S. 519, 534 (1966) stating that "The basic quid pro quo contemplated by the Constitution and the Congress for granting a patent monopoly is the benefit derived by the public from an invention with substantial utility." In explaining the "substantial utility" standard, M.P.E.P. 2107.01 cautions, however, that Office personnel must be careful not to interpret the phrase "immediate benefit to the public" or similar formulations used in certain court decisions to mean that products or services based on the claimed invention must be "currently available" to the public in order to satisfy the utility requirement. "Rather, any reasonable use that an applicant has identified for the invention that can be viewed as providing a public benefit should be accepted as sufficient, at least with regard to defining a "substantial" utility." (M.P.E.P. 2107.01, emphasis added.) Indeed, the Guidelines for Examination of Applications for Compliance With the Utility Requirement, set forth in M.P.E.P. 2107 II (B) (1) gives the following instruction to patent examiners: "If the applicant has asserted that the claimed invention is useful for any particular practical purpose . . . and the assertion would be considered credible by a person of ordinary skill in the art, do not impose a rejection based on lack of utility."

Finally, the Utility Guidelines restate the Patent Office's long established position that any asserted utility has to be "credible." "Credibility is assessed from the perspective of one of ordinary skill in the art in view of the disclosure and any other evidence of record . . . that is probative of the applicant's assertions." (M.P.E.P. 2107 II (B) (1) (ii)) Such standard is presumptively satisfied unless the logic underlying the assertion is seriously flawed, or if the facts upon which the assertion is based are inconsistent with the logic underlying the assertion (Revised Interim Utility Guidelines Training Materials, 1999).

Proper Application of the Legal Standard

Applicants submit that the microarray data provided in the present application is sufficient to establish a credible, specific and substantial utility for PRO3301 polypeptides identified as SEQ ID NO:4. Furthermore, it is Applicants belief that the "universal normal control" described in EXAMPLE 13, is a reliable negative control and is predictive of normal expression levels found in healthy tissues of the same tissue type as the cancerous tumors examined. As such, microarray studies utilizing a "universal normal control" serve as a predictive indicator of overexpression or upregulation of gene expression in cancerous tumors. Applicants would like to draw the Examiner's attention to the fact that parallel studies were concurrently conducted, wherein the microarray analysis of the test sample was compared to expression levels exhibited in a control sample using normal healthy cells representative of the same tissue type as the cancerous samples examined. In this regard, these parallel studies compared the expression levels of PRO3301 polypeptide-encoding nucleic acid sequences in: (1) non-small cell lung adenocarcinoma versus normal lung tissue controls, (2) lung squamous cell carcinoma versus normal lung tissue controls, and (3) colorectal adenocarcinoma versus normal colorectal tissue controls. These contemporaneous studies demonstrated that compared to their normal healthy tissue counterparts, PRO3301 polypeptide-encoding nucleic acid sequences showed, on average, a 2.4 fold, 9.2 fold, and 5.3 fold increase in expression levels in non-small cell lung adenocarcinomas, lung squamous cell carcinomas, and colorectal adenocarcinomas, respectively. In light of the arguments and discussion above, PRO3301 polypeptides of the present invention are significantly overexpressed in specific tumor tissue(s)

when compared to normal controls. Thus, PRO3301 polypeptide-encoding nucleic acid sequences of the present invention can be useful as diagnostic markers for non-small cell lung adenocarcinomas, lung squamous cell carcinomas and colorectal adenocarcinomas. In view of these findings, the novel PRO19598 receptor which binds to a functional PRO3301 (diagnostic marker) would likewise possess a credible, specific and substantial utility. The binding event in and of itself can serve as a useful *in vitro* diagnostic tool to detect and measure the levels of PRO3301 polypeptides in an unknown test sample wherein detected overexpression levels would be suggestive of the presence of lung and colorectal tumors. It is, of course, true that further research would be needed to develop a diagnostic product. However, the fact that such follow-up tests might be necessary, cannot properly lead to the legal conclusion that either PRO3301 or PRO19598 polypeptides or their corresponding encoding nucleic acids lack patentable utility.

As set forth in M.P.E.P., 2107 II (B) (1), if the Applicant has asserted that the claimed invention is useful for any particular purpose, and the assertion would be considered credible by a person of ordinary skill in the art, a rejection based on lack of utility should not be imposed. Accordingly, the Examiner is respectfully requested to reconsider and withdraw the present rejection under 35 U.S.C. §101.

Objections and Rejections under 35 U.S.C. §112, First Paragraph

(1) Claims 22-41 stand rejected under 35 U.S.C. §112, first paragraph as allegedly not being supported by either a credible, specific and substantial asserted utility or a well established utility so that one skilled in the art would not know how to use the claimed invention. Without necessarily agreeing with the propriety of the rejection, Applicants have herein canceled Claims 1-52 rendering the rejection of Claims 22-41 under 35 U.S.C. §112, first paragraph in the instant Office Action moot. Accordingly, Applicants respectfully request that the outstanding rejection of Claims 22-41 under 35 U.S.C. §112, first paragraph be withdrawn.

(2) Claims 1, 5, 7 and 9-17 stand rejected under 35 U.S.C. §112, first paragraph, as allegedly, containing subject matter which was not described in the specification in such a way

as to reasonably convey to one skilled in the relevant art that the inventor(s), at the time the application was filed, had possession of the claimed invention. The Examiner further states "...only isolated polypeptides comprising the amino acid sequence set forth in SEQ ID NO:2, but not the full breadth of the claim meets the written description provision of 35 U.S.C. §112, first paragraph". Accordingly, the Examiner alleges that the specification does not provide adequate written description of the claimed genus. Applicants respectfully traverse.

In light of the amendment to the claims and the discussion above traversing the rejections under 35 U.S.C. §101, Applicants respectfully submit that the specification discloses a credible, specific and substantial utility for both PRO3301 and PRO19598 polypeptides as useful diagnostic tools for detection of lung and colorectal tumors. Applicants have herein amended the rejected genus claims to recite variant nucleic acid sequences encoding a PRO19598 polypeptide, wherein said encoded polypeptide is a receptor for and binds to the ligand polypeptide shown as SEQ ID NO:4. Accordingly, the genus claims, as amended, recite a specific utility for the claimed variant nucleic acids (i.e. said polypeptide serving as a specific receptor that binds to a functional ligand polypeptide). Since the claims are now drawn to a genus defined by both sequence and functional identity, one of ordinary skill in the art would know at the effective priority date of this application, that the Applicants were in possession of the claimed sequences. Accordingly, the Examiner is respectfully requested to reconsider and withdraw the present rejection of the claims under 35 U.S.C. §112, first paragraph.

In light of the above amendments and remarks, Applicants believe that this application is now in condition for immediate allowance and respectfully request that the outstanding objections be withdrawn and this case passed to issue.

The Examiner is invited to contact the undersigned at (650) 225-4563 if any issues may be resolved in that manner.

Attached hereto is a marked-up version of the changes made to the claims by the current amendment. The attached page is captioned "Version with markings to show changes made."

Serial Number: 09/964,994
Filed: September 26, 2001

Respectfully submitted,
GENENTECH, INC.

Date: May 7, 2003

By: Elizabeth M. Barnes
Elizabeth M. Barnes, Ph.D.
Reg. No. 35, 059
Telephone: (650) 225-4563



09157

PATENT TRADEMARK OFFICE

Serial Number: 09/964,994

Filed: September 26, 2001

Version with markings to show changes made

In the Claims

Claims 1-52 have been cancelled

New Claims 53-69 have been added



IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re Application of Audrey Goddard et al. Serial No.: 09/964,994 Filed: September 26, 2001 Title: NOVEL POLYPEPTIDES HAVING SEQUENCE SIMILARITY TO CYTOKINE RECEPTORS AND NUCLEIC ACIDS ENCODING THE SAME	Group Art Unit: 1646 Examiner: KEMMERER, ELIZABETH Confirmation No: 2989 Customer No: 09157 CERTIFICATE OF MAILING I hereby certify that this correspondence is being deposited with the United States Postal Service with sufficient postage as first class mail in an envelope addressed to: Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450 on September 2, 2003 <u>P. Tobin</u> Patty Tobin
--	---

AMENDMENT UNDER 37 C.F.R. §1.111

Commissioner for Patents
P.O. Box 1450
Alexandria, VA 22313-1450

Sir:

This document is responsive to the Final Office Action mailed July 1, 2003 (Paper No. 15) for which a three-month period for response was given. This response is timely filed and the fee therefor. In view of the discussion provided, reconsideration and allowance are respectfully requested.

The Commissioner is authorized to charge any fees which may be required, including extension fees, or credit any overpayment to Deposit Account No. 07-0630.

Amendments to the Claims are reflected in the listing of claims which begins on page two (2) of this paper.

Remarks begin on page seven (7) of this paper.

Amendments to the Claims:

This listing of claims will replace all prior versions, and listings, of claims in the application:

Listing of Claims:

Claims 1-52 have been cancelled.

53. (Currently amended) An isolated nucleic acid molecule having at least 80% sequence identity to:
- (a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;
 - (b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide;
 - (c) the nucleic acid sequence shown as SEQ ID NO:1;
 - (d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or
 - (e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532; wherein ~~said encoded polypeptide~~ said isolated nucleic acid encodes a polypeptide which is a receptor for and binds to the ligand polypeptide shown as SEQ ID NO:4.

54. (Currently amended) An isolated nucleic acid molecule having at least 85% sequence identity to:
- (a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;
 - (b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide;
 - (c) the nucleic acid sequence shown as SEQ ID NO:1;
 - (d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or

(e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532; wherein ~~said encoded polypeptide~~ said isolated nucleic acid encodes a polypeptide which is a receptor for and binds to the ligand polypeptide shown as SEQ ID NO:4.

55. (Currently amended) An isolated nucleic acid molecule having at least 90% sequence identity to:

(a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;

(b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide;

(c) the nucleic acid sequence shown as SEQ ID NO:1;

(d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or

(e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532; wherein ~~said encoded polypeptide~~ said isolated nucleic acid encodes a polypeptide which is a receptor for and binds to the ligand polypeptide shown as SEQ ID NO:4.

56. (Currently amended) An isolated nucleic acid molecule having at least 95% sequence identity to:

(a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;

(b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide;

(c) the nucleic acid sequence shown as SEQ ID NO:1;

(d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or

(e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532; wherein ~~said encoded polypeptide~~ said isolated nucleic acid encodes a polypeptide which is a receptor for and binds to the ligand polypeptide shown as SEQ ID NO:4.

57. (Currently amended) An isolated nucleic acid molecule having at least 99% sequence identity to:

(a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;

(b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide;

(c) the nucleic acid sequence shown as SEQ ID NO:1;

(d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or

(e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532; wherein ~~said encoded polypeptide~~ said isolated nucleic acid encodes a polypeptide which is a receptor for and binds to the ligand polypeptide shown as SEQ ID NO:4.

58. (Previously presented) An isolated nucleic acid molecule comprising:

(a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;

(b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide;

(c) the nucleic acid sequence shown as SEQ ID NO:1;

(d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or

(e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532.

59. (Previously presented) The isolated nucleic acid molecule of Claim 58 comprising a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2.

60. (Previously presented) The isolated nucleic acid molecule of Claim 58 comprising a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide.

61. (Previously presented) The isolated nucleic acid molecule of Claim 58 comprising the nucleic acid sequence shown as SEQ ID NO:1.

62. (Previously presented) The isolated nucleic acid molecule of Claim 58 comprising the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1.

63. (Previously presented) The isolated nucleic acid molecule of Claim 58 comprising the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532.

+64. (Currently amended) An isolated nucleic acid molecule that hybridizes to:

(a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;

(b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide;

(c) the nucleic acid sequence shown as SEQ ID NO:1;

(d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or

(e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532; wherein said isolated nucleic acid encodes a polypeptide

which is a receptor for and binds to the ligand polypeptide shown as SEQ ID NO:4..

65. (Previously presented) The isolated nucleic acid molecule of Claim 64, wherein said hybridization occurs under stringent conditions of 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5xDenhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42 C, with washes at 42 C in 0.2 x SSC (sodium chloride/sodium citrate) and 50% formamide at 55 C, followed by a high-stringency wash consisting of 0.1 x SSC containing EDTA at 55 C.

66. (Currently amended) A vector comprising the isolated nucleic acid molecule of Claim ~~1~~53.

67. (Currently amended) The vector of Claim 66, wherein said isolated nucleic acid is operably linked to control sequences recognized by a host cell transformed with the vector.

68. (Previously presented) A host cell comprising the vector of Claim 66.

69. (Previously presented) The host cell of Claim 68, wherein said cell is a CHO cell, an E. coli or a yeast cell.

REMARKS

Claims 53-69 remain in this application. Claims 1-52 have been previously canceled. Applicants have herein amended Claims 53-57, 64, and 66-67 so as to more clearly define the subject matter claimed therein. In that the newly amended

claims do not introduce new subject matter and are supported in the specification and the claims as originally filed, their entry is respectfully requested.

Objections and Rejections Under 35 U.S.C. § 112, Second Paragraph

Claims 53-57 and 66-69 stand rejected under 35 U.S.C. § 112, second paragraph, as allegedly being indefinite for failing to particularly point out and distinctly claim the subject matter which Applicants regard as the invention. Claims 53-57 have been amended to correct an improper antecedent basis for the phrase "said encoded polypeptide" by deleting the phrase and replacing it with the phrase "said isolated nucleic acid encodes a polypeptide which" as suggested by the Examiner. Claim 66 has been amended to depend from Claim 53 rather than from cancelled Claim 1. In light of the amendments to the claims and the discussion above, Applicants respectively request reconsideration and withdrawal of the outstanding rejections under 35 U.S.C. § 112, second paragraph.

Rejections Under 35 U.S.C. § 101

Claims 53-69 stand rejected under 35 U.S.C. § 101 for allegedly lacking a credible, specific and substantial asserted utility or a well established utility. The Examiner has stated that it is unclear from Example 13 of the specification that the "universal normal control" is an appropriate control. However, the Examiner has indicated that the new data from parallel and concurrent studies that was submitted by Applicant in their response to the January 17, 2003 Office Action may be probative of the utility of the claimed invention. Yet, such new data is more properly presented in the form of a declaration under 37 C.F.R. § 1.132. Applicants herein provide the Declaration of Thomas D. Wu, M.D., Ph.D., wherein Dr. Thomas D. Wu declares that in addition to using the above described "universal normal control" in these microarray studies, parallel studies were concurrently conducted. In these contemporaneous studies, the level of expression of the PRO3301-encoding nucleic acid sequence was compared in both tumor samples and in control samples of the same tissue type as the tumor samples

examined. These concurrent studies demonstrated that compared to their normal healthy tissue counterparts, PRO3301-encoding nucleic acid sequences showed, on average, a 2.4 fold, 9.2 fold, and 5.3 fold increase in expression levels in non-small cell lung adenocarcinomas, lung squamous cell carcinomas, and colorectal adenocarcinomas, respectively. These results demonstrate that PRO3301-encoding nucleic acid sequences are significantly overexpressed in specific tumor tissue(s) when compared to their specific normal tissue controls and thus can be useful as a diagnostic marker for the presence or absence of such specific tumor(s) in a tissue sample of unknown pathology. As such, the PRO19598 polypeptides encoded by the claimed polynucleotides have patentable utility in their being able to bind/label PRO3301 in cancer diagnostic assays.

Applicants respectfully request the acceptance into evidence Dr. Thomas D. Wu's Declaration submitted herein under 37 C.F.R. § 1.132. In light of this declaration, amendment to the claims and the discussion above, Applicants respectfully request reconsideration and withdrawal of the outstanding rejections under 35 U.S.C. § 101.

Rejections Under 35 U.S.C. § 112, First Paragraph

Claims 53-69 are rejected under 35 U.S.C. § 112, first paragraph for allegedly lacking

a credible, specific and substantial asserted utility or a well established utility for the same reasons outlined in the rejections under 35 U.S.C. § 101. Applicants have herein provided the Declaration of Dr. Thomas D. Wu under 37 C.F.R. § 1.132 to provide evidence and support for Applicants' assertion that the claimed invention is supported by a credible, specific and substantial asserted utility or a well established utility so that one skilled in the art would know how to use the claimed invention. In the light of this declaration and the arguments and discussion traversing the rejections under 35 U.S.C. § 101, Applicants respectfully submit that specification discloses a credible, specific and substantial utility for the claimed invention as useful diagnostic tools for detection of lung and colorectal tumors.

In addition, Claims 64-69 are rejected under 35 U.S.C. § 112, first paragraph for allegedly reciting a genus of polynucleotides without indicating functional limitations. The Examiner has stated that Claims 53-63, as previously amended recite structural and functional limitations for the encoded polypeptides resulting in an adequate written description of a claimed genus. Applicants have herein amended Claims 64 and 66-67 to provide the functional limitation "wherein said isolated nucleic acid encodes a polypeptide which is a receptor for and binds to the ligand polypeptide shown as SEQ ID NO:4" in order to provide an adequate written description of the claimed genus. Thus, the genus claims, as amended, recite a specific utility for the claimed variant nucleic acids which encode a polypeptide which is receptor for and binds to a functional ligand polypeptide. Since the claims are now drawn to a genus defined by both sequence and functional identity, one of ordinary skill in the art would know at the effective priority date of this application, that the Applicants were in possession of the claimed sequences.

In view of the Declaration of Dr. Thomas D. Wu submitted herewith, amendments to the claims and the discussion above traversing the rejections under 35 U.S.C. §§ 101 and 112, Applicants respectfully submit that the specification discloses a credible, specific and substantial utility for the claimed invention as useful diagnostic tools for detection of lung and colorectal tumors. Accordingly, the Examiner is respectfully requested to reconsider and withdraw the rejections of the claims under 35 U.S.C. § 112, first paragraph.

In light of the above amendments and remarks, Applicants believe that this application is now in condition for immediate allowance and respectfully request that the outstanding rejections be withdrawn and this case passed to issue.


The Examiner is invited to contact the undersigned at (650) 225-4563 if any issues may be resolved in that manner.

Respectfully submitted,

GENENTECH, INC.

Date: September 2, 2003

By:


Elizabeth M. Barnes, Ph.D.
Reg. No. 35,059

1 DNA Way, MS 49
South San Francisco, CA 94080
Phone: (650) 225-4563
Fax: (650) 952-9881

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

☒ **BLACK BORDERS**

☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**

☐ **FADED TEXT OR DRAWING**

☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**

☐ **SKEWED/SLANTED IMAGES**

☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**

☐ **GRAY SCALE DOCUMENTS**

☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**

☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**

☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.

(12) **United States Patent**
Goddard et al.

(10) **Patent No.:** **US 6,740,520 B2**

(45) **Date of Patent:** **May 25, 2004**

(54) **CYTOKINE RECEPTOR AND NUCLEIC ACIDS ENCODING THE SAME**

(75) **Inventors:** **Audrey Goddard**, San Francisco, CA (US); **Paul J. Godowski**, Hillsborough, CA (US); **Austin L. Gurney**, Belmont, CA (US); **Colin K. Watanabe**, Moraga, CA (US); **William I. Wood**, Hillsborough, CA (US)

(73) **Assignee:** **Genentech, Inc.**, South San Francisco, CA (US)

(*) **Notice:** Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) **Appl. No.:** **09/964,994**

(22) **Filed:** **Sep. 26, 2001**

(65) **Prior Publication Data**

US 2002/0137909 A1 Sep. 26, 2002

Related U.S. Application Data

(63) Continuation-in-part of application No. 09/941,992, filed on Aug. 28, 2001.

(60) Provisional application No. 60/191,015, filed on Mar. 21, 2000.

(30) **Foreign Application Priority Data**

Mar. 30, 2000 (WO) PCT/US00/08439
 Feb. 28, 2001 (WO) PCT/US01/06520

(51) **Int. Cl.⁷** **C12N 5/00**

(52) **U.S. Cl.** **435/325; 435/252.3; 435/254.11; 435/320.1; 536/23.1; 536/23.5**

(58) **Field of Search** **536/231, 23.5; 435/69.1, 320.1, 325, 252.3, 254.11**

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,179,337 A 12/1979 Davis et al.
 4,275,149 A 6/1981 Litman et al.
 4,301,144 A 11/1981 Iwashita et al.
 4,496,689 A 1/1985 Mitra
 4,640,835 A 2/1987 Shimizu et al.
 4,657,760 A 4/1987 Kung et al.
 4,670,417 A 6/1987 Iwasaki et al.
 4,676,980 A 6/1987 Segal et al.
 4,736,866 A 4/1988 Leder et al.
 4,791,192 A 12/1988 Nakagawa et al.
 4,816,567 A 3/1989 Cabilly et al.
 4,870,009 A 9/1989 Evans et al.
 5,206,344 A 4/1993 Katre et al.
 5,225,212 A 7/1993 Martin et al.
 5,364,934 A 11/1994 Drayna et al.
 5,536,637 A 7/1996 Jacobs
 5,545,806 A 8/1996 Lonberg et al.
 5,545,807 A 8/1996 Surani et al.
 5,569,825 A 10/1996 Lonberg et al.
 5,625,126 A 4/1997 Lonberg et al.
 5,633,425 A 5/1997 Lonberg et al.
 5,654,010 A 8/1997 Johnson et al.
 5,661,016 A 8/1997 Lonberg et al.

OTHER PUBLICATIONS

Kotenko et al., 2001, J. Immunol. 166:7096-7103.*

Altschul and Gish, "Local Alignment Statistics" *Methods in Enzymology* 266:460-480 (1996).

Altschul, S.F. et al., "Gapped Blast and PSI-Blast: a new generation of protein database search programs" *Nucleic Acids Research* 25:3389-3402 (1997).

Anderson, W.F., "Human gene therapy" *Science* 256(5058):808-813 (May 8, 1992).

Beal and Dervan, "Second Structural Motif for Recognition of DNA by Oligonucleotide-Directed Triple-Helix Formation" *Science* 251:1360-1363 (1991).

Boerner et al., "Production of Antigen-Specific Human Monoclonal Antibodies From In Vitro-Primed Human Splenocytes" *The Journal of Immunology* 147(1):86-95 (1991).

Bradley, "Production and Analysis of Chimaeric Mice" *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E. J. Robertson, ed., IRL, Oxford, Chapter 5, pp. 113-151 (1987).

Brennan et al., "Preparation of bispecific antibodies by chemical recombination of monoclonal immunoglobulin G₁ fragments" *Science* 229:81-83 (Jul. 1985).

Brodeur et al., "Mouse-Human Myeloma Partners for the Production of Heterohybridomas" *Monoclonal Antibody Production Techniques and Application*, New York:Marcel Dekker, Inc. pp. 51-63 (1987).

Caron et al., "Engineered humanized dimeric forms of IgG are more effective antibodies" *Journal of Experimental Medicine* 176(4):1191-1195 (1992).

Carter et al., "Improved Oligonucleotide Site-Directed Mutagenesis Using M13 Vectors" *Nucl. Acids Res.* 13(12):4431-4443 (Jun. 25, 1985).

Chevray and Nathans, "Protein Interaction Cloning in Yeast: Identification of Mammalian Proteins That React with the Leucine Zipper of Jun" *Proc. Natl. Acad. Sci. USA* 89:5789-5793 (Jul. 1992).

Chien et al., "The Two-Hybrid System: A method to identify and clone genes for proteins that interact with a protein of interest" *Proc. Natl. Acad. Sci. USA* 88:9578-9582 (1991).

Chothia, "The Nature of the Accessible and Buried Surfaces in Proteins" *Journal Mol. Biol.* 105:1-14 (1976).

(List continued on next page.)

Primary Examiner—Elizabeth Kemmerer

(74) **Attorney, Agent, or Firm**—Elizabeth M. Barnes

(57) **ABSTRACT**

The present invention is directed to novel cytokine receptors having sequence similarity to AF18497_1 and to nucleic acid molecules encoding those cytokine receptors. Also provided herein are vectors and host cells comprising those nucleic acid sequences, chimeric polypeptide molecules comprising the polypeptides of the present invention fused to heterologous polypeptide sequences, antibodies which bind to the polypeptides of the present invention and to methods for producing the polypeptides of the present invention.

16 Claims, 4 Drawing Sheets

OTHER PUBLICATIONS

- Cleland., "Design and Production of Single Immunization Vaccines Using Polylactide Polyglycolide Microsphere Systems." *Vaccine Design: The Subunit and Adjuvant Approach*, Powell and Newman, New York:Plenum Press pp. 439-462 (1995).
- Cohen, Jack S. ed. *Oligodeoxynucleotides: Antisense Inhibitors of Gene Expression*, Boca Raton, FL: CRC Press, Inc. (1989).
- Cole et al., "The EBV-Hybridoma Technique and Its Application to Human Lung Cancer" *Monoclonal Antibodies and Cancer Therapy*, New York: Alan R. Liss, Inc. pp. 77-96 (1985).
- Coligan et al. *Current Protocols in Immunology*, Chapter 5, 1(2) (1991).
- Cooney et al., "Site-Specific Oligonucleotide Binding Represses Transcription of the Human c-myc Gene in Vitro" *Science* 241:456-459 (Jul. 22, 1988).
- Creighton, Thomas E. *Proteins: Structures and Molecular Properties*, N.Y.: W.H. Freeman & Co. (1983).
- David and Reisfeld., "Protein Iodination with Solid State Lactoperoxidase." *Biochemistry* 13(5):1014-1021 (1974).
- Dzau et al., "Gene therapy for cardiovascular disease" *Trends in Biotechnology* 11:205-210 (1993).
- Evan et al., "Isolation of Monoclonal Antibodies Specific for Human c-myc Proto-Oncogene Product" *Molecular & Cellular Biology* 5:3610-3616 (1985).
- Field et al., "Purification of a RAS-Responsive Adenyl Cyclase Complex from *Saccharomyces cerevisiae* by Use of an Epitope Addition Method" *Molecular & Cellular Biology* 8:2159-2165 (1988).
- Fields and Song, "A novel genetic system to detect protein-protein interactions" *Nature* 340:245-246 (1989).
- Fishwild et al., "High-Avidity Human IgG kappa Monoclonal Antibodies From a Novel Strain of Minilocus Transgenic Mice" *Nature Biotechnology* 14(7):845-851 (Jul. 1996).
- Gruber et al., "Efficient Tumor Cell Lysis Mediated by a Bispecific Single Chain Antibody Expressed in *Escherichia Coli*" *Journal of Immunology* 152:5368-5374 (1994).
- Holliger et al., "Diabodies": Small Bivalent and Bispecific Antibody Fragments," *Proc. Natl. Acad. Sci. USA* 90:6444-48 (Jul. 1993).
- Holmes et al., "Structure and Functional Expression of a Human Interleukin-8 Receptor" *Science* 253(5025):1278-1280 (1991).
- Hoogenboom and Winter, "By-Passing Immunisation: Human Antibodies from Synthetic Repertoires of Germline V_H Gene Segments Rearranged in Vitro" *J. Mol. Biol.* 227:381-388 (1992).
- Hopp et al., "A Short Polypeptide Marker Sequence Useful for Recombinant Protein Identification and Purification" *Bio/Technology* 6:1204-1210 (1988).
- Hora et al., "Controlled Release of Interleukin-2 From Biodegradable Microspheres" *Bio/Technology* 8:755-758 (Aug. 1990).
- Johnson et al., "A Month-Long Effect From a Single Injection of Microencapsulated Human Growth Hormone" *Nature Medicine* 2:795-799 (1996).
- Jones, P.T. et al., "Replacing the Complementarity-determining Regions in a Human Antibody with Those From a Mouse" *Nature* 321:522-525 (May 29, 1986).
- Klein et al., "Selection for Genes Encoding Secreted Proteins and Receptors" *Proc. Natl. Acad. Sci. USA* 93(14):7108-7113 (1996).
- Kohler, G. and Milstein, C., "Continuous Cultures of Fused Cells Secreting Antibody of Predefined Specificity" *Nature* 256:495-497 (Aug. 7, 1975).
- Kosteiny et al., "Formation of a Bispecific Antibody by the Use of Leucine Zippers" *Journal of Immunology* 148(5):1547-1553 (1992).
- Kozbor et al., "A Human Hybrid Myeloma for Production of Human Monoclonal Antibodies" *Journal of Immunology* 133(6):3001-3005 (1984).
- Lee et al., "Complexes formed by (pyrimidine)_n (purine)_n DNAs on lowering the pH are three-stranded" *Nucl. Acids Res.* 6:3073-3091 (1979).
- Lewis, "Controlled release of bioactive agents from lactide/glycolide polymers" *Biodegradable Polymers as Drug Delivery Systems*, M. Chasin and R Langer, New York: Marcel Dekker pp. 1-41 (1990).
- Li et al., "Targeted mutation of the DNA methyltransferase gene results in embryonic lethality" *Cell* 69:915-926 (1992).
- Lonberg and Huszar, "Human Antibodies From Transgenic Mice." *International Reviews of Immunology* 13(1):65-93 (1995).
- Lonberg et al., "Antigen-specific human antibodies from mice comprising four distinct genetic modifications" *Nature* 368(6474):856-859 (Apr. 28, 1994).
- Lutz-Freyermuth et al., "Quantitative Determination That One of Two Potential RNA-binding Domains of the A Protein Component of the U1 Small Nuclear Ribonucleoprotein Complex Binds with High Affinity to Stem-loop II of U1 RNA" *Proc. Natl. Acad. Sci. USA* 87:6393-6397 (1990).
- Marasco et al., "Design, intracellular expression, and activity of a human anti-human immunodeficiency virus type 1 gp 120 single-chain antibody" *Proc. Natl. Acad. Sci. USA* 90:7889-7893 (1993).
- Marks et al., "By-Passing Immunization: Building High Affinity Human Antibodies by Chain Shuffling" *Bio/Technology* 10:779-783 (1992).
- Marks et al., "By-Passing Immunization: Human Antibodies From V-gene Libraries Displayed On Phage." *J. Mol. Biol.* 222:581-597 (1991).
- Martin et al., "GAP Domains Responsible for Ras p21-Dependent Inhibition of Muscarinic Atrial K⁺ Channel Currents" *Science* 255:192-194 (1992).
- Merrifield, R.B., "Solid Phase Peptide Synthesis. I. The Synthesis of a Tetrapeptide" *J. Am. Chem. Soc.* 85:2149-2154 (1963).
- Milstein and Cuello, "Hybrid Hybridomas and Their Use in Immunohistochemistry" *Nature* 305:537-540 (Oct. 1983).
- Mordenti et al., "The Use of Interspecies Scaling in Toxicokinetics" *Toxicokinetics and New Drug Development*, Yacobi et al., eds., New York: Pergamon Press pp. 42-96 (1989).
- Morrison, S., "Immunology: Success in specification" *Nature* 368(6474):812-813 (Apr. 28, 1994).
- Munson and Rodbard, "Ligand: A Versatile Computerized Approach for characterization of Ligand-Binding Systems" *Analytical Biochemistry* 107:220-239 (1980).
- Neuberger, M., "Generating high-avidity human Mabs in mice" *Nature Biotechnology* 14(7):826 (Jul. 1996).

- Nielsen et al., "Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites" *Protein Engineering* 10(1):1-6 (1997).
- Nygren, H., "Conjugation of Horseradish Peroxidase to Fab Fragments with Different Homobifunctional and Heterobifunctional Cross-Linking Reagents" *The Journal of Histochemistry and Cytochemistry* 30(5):407-412 (1982).
- Okano et al., "Myelin Basic Protein Gene and the Function of Antisense RNA in Its Repression in Myelin-Deficient Mutant Mouse" *J. of Neurochemistry* 56(2):560-567 (1991).
- Paborsky et al., "Mammalian Cell Transient Expression of Tissue Factor for the Production of Antigen" *Protein Eng.* 3(6):547-553 (1990).
- Pain et al., "Preparation of Protein λ -Peroxidase Monoconjugate Using a Heterobifunctional Reagent, and its Use in Enzyme Immunoassays" *Journal of Immunological Methods* 40:219-230 (1981).
- Pluckthun, A., "Antibodies from *Escherichia coli*" *The Pharmacology of Monoclonal Antibodies*, Rosenberg and Moore, New York:Springer-Verlag, Chapter 11, vol. 113:269-315 (1994).
- Presta, L., "Antibody Engineering" *Curr. Op. Struct. Biol.* 2:593-596 (1992).
- Riechmann, L. et al., "Reshaping Human Antibodies for Therapy" *Nature* 332:323-327 (Mar. 24, 1988).
- Rossi, J.J., "Making ribozymes work in cells" *Current Biology* 4:469-471 (1994).
- Sambrook et al. *Molecular Cloning: A Laboratory Manual*, Second edition, New York: Cold Spring Harbor Laboratory Press (1989).
- Shalaby et al., "Development of Humanized Bispecific Antibodies Reactive with Cytotoxic Lymphocytes and Tumor Cells Overexpressing the HER2 Protooncogene" *Journal of Experimental Medicine* 175:217-225 (Jan. 1, 1992).
- Shopes, "A genetically engineered human IgG mutant with enhanced cytolytic activity" *Journal of Immunology* 148(9):2918-2922 (1992).
- Skinner et al., "Use of the Glu-Glu-Phe C-terminal Epitope for Rapid Purification of the Catalytic Domain of Normal and Mutant ras GTPase-activating Proteins" *Journal of Biological Chemistry* 266:14163-14166 (1991).
- Stein and Cohen., "Oligodeoxynucleotides as Inhibitors of Gene Expression: A Review." *Cancer Research* 48:2659-2668 (May 15, 1988).
- Stevenson et al., "A Chimeric Antibody with Dual Fc Regions (bisFabFc) Prepared by Manipulations at the IgG Hinge" *Anti-Cancer Drug Design* 3(4):219-230 (1989).
- Stewart et al. *Solid-phase Peptides Synthesis*, San Francisco, CA: W.H. Freeman Co. (1969).
- Suresh et al., "Bispecific Monoclonal Antibodies from Hybrid Hybridomas" *Methods in Enzymology* 121:210-228 (1986).
- Thomas and Capecchi, "Site-Directed Mutagenesis by Gene Targeting in Mouse Embryo-Derived Stem Cells" *Cell* 51:503-512 (1987).
- Thomas, P., "Hybridization of Denatured RNA and Small DNA Fragments Transferred to Nitrocellulose" *Proc. Natl. Acad. Sci. USA* 77(9):5201-5205 (Sep. 1980).
- Traunecker et al., "Bispecific Single Chain Molecules (Janusins) Target Cytotoxic Lymphocytes on HIV Infected Cells" *EMBO Journal* 10(12):3655-3659 (1991).
- Tutt et al., "Trispecific F(ab')₃ Derivatives That Use Cooperative Signaling Via the TCR/CD3 Complex and CD2 to Activate and Redirect Resting Cytotoxic T Cells" *J. Immunol.* 147(1):60-69 (1991).
- van der Krol, Mol and Stuitje., "Modulation of Eukaryotic Gene Expression by Complementary RNA or DNA Sequences." *Biotechniques*. 6(10):958-976 (1998).
- Verhoeyan, M. et al., "Reshaping Human Antibodies: Grafting an Antilysozyme Activity" *Science* 239:1534-1536 (Mar. 25, 1988).
- von Heijne et al., "A New Method for Predicting Signal Sequence Cleavage Sites." *Nucl. Acids Res.* 14(11):4683-4690 (1986).
- Wagner et al., "Transferrin-polycation conjugates as carriers for DNA uptake into cells" *Proc. Natl. Acad. Sci.* 87:3410-3414 (1990).
- Wells et al., "Cassette Mutagenesis: an Efficient Method for Generation of Multiple Mutations at Defined Sites" *Gene* 34(2-3):315-323 (1985).
- Wells et al., "Importance of Hydrogen-Bond Formation in Stabilizing the Transition State of Subtilisin" *Philos. Trans. Royal Soc. London Ser. A* 317:415-423 (1986).
- Wolff et al., "Monoclonal antibody homodimers: enhanced antitumor activity in nude mice" *Cancer Research* 53(11):2560-2565 (1993).
- Wu et al., "Receptor-mediated in vitro gene transformation by a soluble DNA carrier system" *Journal of Biological Chemistry* 262(10):4429-4432 (1987).
- Zamecnik et al., "Inhibition of replication and expression of human T-cell lymphotropic virus type III in cultured cells by exogenous synthetic oligonucleotides complementary to viral RNA" *Proc. Natl. Acad. Sci.* 83:4143-4146 (1986).
- Zapata et al., "Engineering linear F(ab')₂ fragments for efficient production in *Escherichia coli* and enhanced antiproliferative activity" *Protein Engineering* 8(10):1057-1062 (1995).
- Zola, "Using Monoclonal Antibodies: Soluble Antigens" *Monoclonal Antibodies: A Manual of Techniques*, CRC Press, Chapter 6, pp. 147-158 (1987).
- Zoller and Smith, "Oligonucleotide-directed Mutagenesis Using M13-derived Vectors: An Efficient and General Procedure for the Production of Point Mutations in Any Fragment of DNA" *Nucl. Acids Res.* 10(20):6487-6500 (1982).

FIGURE 1

CAGTTTCTTCATCTGTAAACATCAAATGAATAATAATACCAATCTCCTAGACTTCATAAGA
GGATTAAACAAAGACAAAATATGGGAAAAACATAACATGGCGTCCCATAATTATTAGATCT
TATTATTGACACTAAAATGGCATTAAAATTACCAAAGGAAGACAGCATCTGTTTCCTCT
TTGGTCCTGAGCTGGTTAAAAGGAACACTGGTTGCCTGAACAGTCACACTTGCAACCATG
ATGCCCTAAACATTGCTTTCTAGGCTTCCTCATCAGTTTCTTCCTTACTGGTGTAGCAGGA
ACTCAGTCAACGCATGAGTCTCTGAAGCCTCAGAGGGTACAATTCAGTCCCGAAATTTT
CACAACATTTTGCAATGGCAGCCTGGGAGGGCACTTACTGGCAACAGCAGTGTCTATTTT
GTGCAGTACAAAATCATGTTCTCATGCAGCATGAAAAGCTCTCACCAGAAGCCAAGTGGA
TGCTGGCAGCACATTTCTTGTAACCTCCCAGGCTGCAGAACATTGGCTAAATATGGACAG
AGACAATGGAAAAATAAAGAAGACTGTTGGGGTACTCAAGAACTCTCTTGTGACCTTACC
AGTGAAACCTCAGACATACAGGAACCTTATTACGGGAGGGTGAGGGCGGCTCGGCTGGG
AGCTACTCAGAATGGAGCATGACGCCGCGGTTCACTCCCTGGTGGGAAACAAAAATAGAT
CCTCCAGTCATGAATATAACCCAAGTCAATGGCTCTTTGTTGGTAATTCTCCATGCTCCA
AATTTACCATATAGATACCAAAGGAAAAAATGTATCTATAGAAGATTACTATGAACTA
CTATACCGAGTTTTTATAATTAACAATTCAGTAGAAAAGGAGCAAAGGTTTATGAAGGG
GCTCACAGAGCGGTTGAAATTGAAGCTCTAACACCACACTCCAGCTACTGTGTAGTGGCT
GAAATATATCAGCCCATGTTAGACAGAAGAAGTCAGAGAAGTGAAGAGAGATGTGTGGAA
ATTCCAT**G**ACTTGTGGAATTTGGCATTTCAGCAATGTGGAAATTCTAAAGCTCCCTGAGAA
CAGGATGACTCGTGTGTTGAAGGATCTTATTTAAAATTGTTTTTGTATTTTCTTAAAGCAA
TATTCAGTGTACACCTTGGGGACTTCTTTGTTTATCCATTCTTTTATCCTTTATATTTT
ATTTGTAAACTATATTTGAACGACATTCCCCCGAAAAATTGAAATGTAAAGATGAGGCA
GAGAATAAAGTGTTCTATGAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

FIGURE 2

```
></usr/seqdb2/sst/DNA/Dnaseqs.min/ss.DNA145887
><subunit 1 of 1, 262 aa, 1 stop
><MW: 30419, pI: 8.44, NX(S/T): 5
MPKHCFLGFLISFFLTGVAGTQSTHESLKPQRVQFQSRNFHNIQWQPGRALTGNSSVYF
VQYKIMFSCSMKSSHQKPSGCWQHISCNFPGCRTLAKYGQRQWKNKEDCWGTQELSCDLT
SETSDIQEPYYGRVRAASAGSYSEWSMTPRFTPWWETKIDPPVMNITQVNGSLLVILHAP
NLPYRYQKEKNVSIEDYYELLYRVFIINNSLEKEQKVYEGAHRAVEIEALTPHSSYCVVA
EIYQPMLDRRSQRSEERCVEIP
```

Important features of the protein:**Signal peptide:**

Amino Acids 1-20

N-glycosylation sites:

Amino acids 55-59;165-169;170-174;191-195;208-212

N-myristoylation sites:

Amino acids 17-23;20-26;220-226

FIGURE 3

TGAAATGACTTCCACGGCTGGGACGGGAACCTTCCACCCACAGCTATGCCTCTGATTGGTGA
ATGGTGAAGGTGCCTGTCTAACTTTTCTGTAAAAAGAACCAGCTGCCTCCAGGCAGCCAGCC
CTCAAGCATCACTTACAGGACCAGAGGGACAAGACATGACTGTGATGAGGAGCTGCTTTCGC
CAATTTAACACCAAGAAGAATTGAGGCTGCTTGGGAGGAAGGCCAGGAGGAACACGAGACTG
AGAGATGAATTTTCAACAGAGGCTGCAAAGCCTGTGGACTTTAGCCAGACCCCTTCTGCCCTC
CTTTGCTGGCGACAGCCTCTCAAATGCAGATGGTTGTGCTCCCTTGCCTGGGTTTACCCCTG
CTTCTCTGGAGCCAGGTATCAGGGGCCAGGGCCAAGAATTCACCTTTGGGCCCTGCCAAGT
GAAGGGGGTTGTTCCCCAGAACTGTGGGAAGCCTTCTGGGCTGTGAAAGACACTATGCAAG
CTCAGGATAACATCACGAGTGCCCGGCTGCTGCAGCAGGAGGTTCTGCAGAACGTCTCGGAT
GCTGAGAGCTGTTACCTTGTCCACACCCTGCTGGAGTTCTACTTGAAAACGTGTTTTCAAAAA
CCACCACAATAGAACAGTTGAAGTCAGGACTCTGAAGTCATTCTCTACTCTGGCCAACAACCT
TTGTTCTCATCGTGTCACAACTGCAACCCAGTCAAGAAAATGAGATGTTTTCCATCAGAGAC
AGTGACACACAGGCGGTTTCTGCTATTCGGGAGAGCATTCAAACAGTTGGACGTAGAAGCAGC
TCTGACCAAAGCCCTTGGGGAAGTGGACATTCTTCTGACCTGGATGCAGAAAATCTACAAGC
TCTGAATGTCTAGACCAGGACCTCCCTCCCCCTGGCACTGGTTTGTTCCTGTGTCATTTCA
AACAGTCTCCCTTCCTATGCTGTTCACTGGACACTTCACGCCCTTGGCCATGGGTCCCATT
TTGGCCCAGGATTATTGTCAAAGAAGTCATTCTTTAAGCAGCGCCAGTGACAGTCAGGGAAG
GTGCCTCTGGATGCTGTGAAGAGTCTACAGAGAAGATTCTTGTATTTATTACAACTCTATTT
AATTAATGTCAGTATTTCAACTGAAGTTCTATTTATTTGTGAGACTGTAAGTTACATGAAGG
CAGCAGAATATTGTGCCCCATGCTTCTTTACCCCTCACAATCCTTGCCACAGTGTGGGGCAG
TGGATGGGTGCTTAGTAAGTACTTAATAAACTGTGGTGCTTTTTTTGGCCTGTCTTTGGATT
GTTAAAAAACAGAGAGGGATGCTTGGATGTAAACTGAACTTCAGAGCATGAAAATCACACT
GTCTTCTGATATCTGCAGGGACAGAGCATTGGGGTGGGGTAAGGTGCATCTGTTTGAAAAG
TAAACGATAAAATGTGGATTAAAGTGCCAGCACAAAGCAGATCCTCAATAAACATTTTATT
TCCCACCCACACTCGCCAGCTCACCCCATCATCCCTTCCCTTGGTGCCCTCCTTTTTTTTTT
TATCCTAGTCATTCTTCCCTAATCTTCCACTTGAGTGTCAAGCTGACCTTGCTGATGGTGAC
ATTGCACCTGGATGTACTATCCAATCTGTGATGACATTCCCTGCTAATAAAAGACAACATAA
CTCCAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

FIGURE 4

></usr/seqdb2/sst/DNA/Dnaseqs.full/ss.DNA88002

><subunit 1 of 1, 206 aa, 1 stop

><MW: 23799, pI: 9.12, NX(S/T): 3

MNFQQRQLQSLWTLARPFPCPLLATASQMOMVVLPCLGFTLLLWSQVSGAQQGEFHFGPCQVK

GVVPQKLWEAFWAVKDTMQAQDNITSARLLQQEVLQNVSDAESCYLEVHTLLEFYLKTVFKNH

HNRTVEVRTLKSFSTLANNFVLIVSQLQPSQENEMFSIRDSAHRRFLLFRAFKQLDVEAAL

TKALGEVDILLTWMQKFYKL

Signal sequence:

amino acids 1-42

N-glycosylation sites.

amino acids 85-89, 99-103, 126-130

CYTOKINE RECEPTOR AND NUCLEIC ACIDS ENCODING THE SAME

This application claims priority under 35 U.S.C. § 119(e) of U.S. provisional Application Serial No. 60/191,015 filed on Mar. 21, 2000, and claims priority under 35 U.S.C. § 119 (a)–(d) of PCT Application Nos. PCT/US00/08439 filed Mar. 30, 2000 and PCT/US01/06520 filed Feb. 28, 2001, and is a continuation-in-part, claiming priority under 35 U.S.C. § 120, of U.S. application Ser. No. 09/941,992 filed Aug. 28, 2001 the entire disclosures of which are hereby expressly incorporated by reference.

FIELD OF THE INVENTION

The present invention relates generally to the identification and isolation of novel DNA and to the recombinant production of novel cytokine receptors, designated herein as “PRO19598” polypeptides.

BACKGROUND OF THE INVENTION

Extracellular proteins play important roles in, among other things, the formation, differentiation and maintenance of multicellular organisms. The fate of many individual cells, e.g., proliferation, migration, differentiation, or interaction with other cells, is typically governed by information received from other cells and/or the immediate environment. This information is often transmitted by secreted polypeptides (for instance, mitogenic factors, survival factors, cytotoxic factors, differentiation factors, neuropeptides, and hormones) which are, in turn, received and interpreted by diverse cell receptors or membrane-bound proteins. These secreted polypeptides or signaling molecules normally pass through the cellular secretory pathway to reach their site of action in the extracellular environment.

Secreted proteins have various industrial applications, including as pharmaceuticals, diagnostics, biosensors and bioreactors. Most protein drugs available at present, such as thrombolytic agents, interferons, interleukins, erythropoietins, colony stimulating factors, and various other cytokines, are secretory proteins. Their receptors, which are membrane proteins, also have potential as therapeutic or diagnostic agents. Efforts are being undertaken by both industry and academia to identify new, native secreted proteins. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel secreted proteins. Examples of screening methods and techniques are described in the literature [see, for example, Klein et al., *Proc. Natl. Acad. Sci.*, 93:7108–7113 (1996); U.S. Pat. No. 5,536,637].

Membrane-bound proteins and receptors can play important roles in, among other things, the formation, differentiation and maintenance of multicellular organisms. The fate of many individual cells, e.g., proliferation, migration, differentiation, or interaction with other cells, is typically governed by information received from other cells and/or the immediate environment. This information is often transmitted by secreted polypeptides (for instance, mitogenic factors, survival factors, cytotoxic factors, differentiation factors, neuropeptides, and hormones) which are, in turn, received and interpreted by diverse cell receptors or membrane-bound proteins. Such membrane-bound proteins and cell receptors include, but are not limited to, cytokine receptors, receptor kinases, receptor phosphatases, receptors involved in cell-cell interactions, and cellular adhesion molecules like selectins and integrins. For instance, transduction of signals that regulate cell growth and differentiation is

regulated in part by phosphorylation of various cellular proteins. Protein tyrosine kinases, enzymes that catalyze that process, can also act as growth factor receptors. Examples include fibroblast growth factor receptor and nerve growth factor receptor.

Membrane-bound proteins and receptor molecules have various industrial applications, including as pharmaceutical and diagnostic agents. Receptor immunoadhesins, for instance, can be employed as therapeutic agents to block receptor-ligand interactions. The membrane-bound proteins can also be employed for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction.

Efforts are being undertaken by both industry and academia to identify new, native receptor or membrane-bound proteins. Many efforts are focused on the screening of mammalian recombinant DNA libraries to identify the coding sequences for novel receptor or membrane-bound proteins.

“Decoy” receptors are well documented in the art. A decoy receptor is generally defined as a receptor that does not transduce an intracellular signal due to lack of a required domain. The decoy receptor may be membrane bound or it may be a soluble secreted form. Decoy receptors represent a unique mechanism for regulating the sensitivity of cells to a particular ligand, as increases in the amount of decoy receptor results in a decrease in the amount of ligand that is available to bind to functional receptors. This titration of ligand by a decoy receptor may modulate the cellular response to the ligand, for example resulting in decreased or increased inflammation, differentiation, proliferation or apoptosis.

We herein describe the identification and characterization of novel cytokine receptors, designated herein as PRO19598 polypeptides.

SUMMARY OF THE INVENTION

A cDNA clone (designated herein as DNA145887-2849) has been identified that encodes a cytokine receptor designated in the present application as “PRO19598”. PRO19598 is a secreted cytokine receptor (otherwise known as a Decoy receptor) which binds to a cytokine in the IL-10 family which is designated herein as a PRO3301 polypeptide ligand (shown in FIG. 4; SEQ ID NO:7). The PRO3301 polypeptide ligand is encoded by a nucleic acid molecule designated herein as DNA88002 (shown in FIG. 3; SEQ ID NO:6).

In one embodiment, the invention provides an isolated nucleic acid molecule comprising a nucleotide sequence that encodes a PRO19598 polypeptide.

In one aspect, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alterna-

tively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule encoding a PRO19598 polypeptide having the sequence of amino acid residues from 1 or 21 to about 262, inclusive, of FIG. 2 (SEQ ID NO:2), or (b) the complement of the nucleotide sequence of (a).

In another aspect, the isolated nucleic acid molecule comprises (a) a nucleotide sequence encoding a PRO19598 polypeptide having the sequence of amino acid residues from 1 or 21 to about 262, inclusive, of FIG. 2 (SEQ ID NO:2), or (b) the complement of the nucleotide sequence of (a).

In other aspects, the isolated nucleic acid molecule comprises a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule having the sequence of nucleotides from 241 or 301 to about 1026, inclusive, of FIG. 1 (SEQ ID NO:1), or (b) the complement of the nucleotide sequence of (a).

In another aspect, the isolated nucleic acid molecule comprises (a) the nucleotide sequence from 241 or 301 to about 1026, inclusive, of FIG. 1 (SEQ ID NO:1), or (b) the complement of the nucleotide sequence of (a).

In a further aspect, the invention concerns an isolated nucleic acid molecule comprising a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least

about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) a DNA molecule that encodes the same mature polypeptide encoded by any of the human protein cDNA deposited with the ATCC on Mar. 21, 2000 under ATCC Deposit No. PTA-1532 (DNA145887-2849), or (b) the complement of the DNA molecule of (a). In a preferred embodiment, the isolated nucleic acid molecule comprises (a) a nucleotide sequence encoding the same mature polypeptide encoded by the human protein cDNA deposited with the ATCC on Mar. 21, 2000 under ATCC Deposit No. PTA-1532 (DNA145887-2849), or (b) the complement of the DNA molecule of (a).

In a another aspect, the invention concerns an isolated nucleic acid molecule comprising a nucleotide sequence having at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) the full-length polypeptide coding sequence of the human protein cDNA deposited with the ATCC on Mar. 21, 2000 under ATCC Deposit No. PTA-1532 (DNA145887-2849), or (b) the complement of the DNA molecule of (a). In a preferred embodiment, the isolated nucleic acid molecule comprises (a) the full-length coding sequence of the human protein cDNA deposited with the ATCC on Mar. 21, 2000 under ATCC Deposit No. PTA-1532 (DNA145887-2849), or (b) the complement of the DNA molecule of (a).

In another aspect, the invention concerns an isolated nucleic acid molecule which encodes an active PRO19598 polypeptide as defined below comprising a nucleotide sequence that hybridizes to the complement of a nucleic acid sequence that encodes amino acids 1 or 21 to about 262, inclusive, of FIG. 2 (SEQ ID NO:2). Preferably, hybridization occurs under stringent hybridization and wash conditions.

In yet another aspect, the invention concerns an isolated nucleic acid molecule which encodes an active PRO19598 polypeptide as defined below comprising a nucleotide sequence that hybridizes to the complement of the nucleic acid sequence between nucleotides 241 or 301 and about 1026, inclusive, of FIG. 1 (SEQ ID NO:1). Preferably, hybridization occurs under stringent hybridization and wash conditions.

In a further aspect, the invention concerns an isolated nucleic acid molecule having at least about 406 nucleotides

and which is produced by hybridizing a test DNA molecule under stringent conditions with (a) a DNA molecule encoding a PRO19598 polypeptide having the sequence of amino acid residues from 1 or 21 to about 262, inclusive, of FIG. 2 (SEQ ID NO:2), or (b) the complement of the DNA molecule of (a), and, if the test DNA molecule has at least about an 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) or (b), and isolating the test DNA molecule.

In another aspect, the invention concerns an isolated nucleic acid molecule comprising (a) a nucleotide sequence encoding a polypeptide scoring at least about 80% positives, alternatively at least about 81% positives, alternatively at least about 82% positives, alternatively at least about 83% positives, alternatively at least about 84% positives, alternatively at least about 85% positives, alternatively at least about 86% positives, alternatively at least about 87% positives, alternatively at least about 88% positives, alternatively at least about 89% positives, alternatively at least about 90% positives, alternatively at least about 91% positives, alternatively at least about 92% positives, alternatively at least about 93% positives, alternatively at least about 94% positives, alternatively at least about 95% positives, alternatively at least about 96% positives, alternatively at least about 97% positives, alternatively at least about 98% positives and alternatively at least about 99% positives when compared with the amino acid sequence of residues 1 or 21 to 262, inclusive, of FIG. 2 (SEQ ID NO:2), or (b) the complement of the nucleotide sequence of (a).

In a specific aspect, the invention provides an isolated nucleic acid molecule comprising DNA encoding a PRO19598 polypeptide without the N-terminal signal sequence and/or the initiating methionine, or is complementary to such encoding nucleic acid molecule. The signal peptide has been tentatively identified as extending from amino acid position 1 to amino acid position 20 in the sequence of FIG. 2 (SEQ ID NO:2). It is noted, however, that the C-terminal boundary of the signal peptide may vary, but most likely by no more than about 5 amino acids on either side of the signal peptide C-terminal boundary as initially identified herein, wherein the C-terminal boundary of the signal peptide may be identified pursuant to criteria routinely employed in the art for identifying that type of amino acid sequence element (e.g., Nielsen et al., *Prot. Eng.* 10:1-6 (1997) and von Heinje et al., *Nucl. Acids. Res.* 14:4683-4690 (1986)). Moreover, it is also recognized that, in some cases, cleavage of a signal sequence from a secreted

polypeptide is not entirely uniform, resulting in more than one secreted species. These polypeptides, and the polynucleotides encoding them, are contemplated by the present invention. As such, for purposes of the present application, the signal peptide of the PRO19598 polypeptide shown in FIG. 2 (SEQ ID NO:2) extends from amino acids 1 to X of FIG. 2 (SEQ ID NO:2), wherein X is any amino acid from 16 to 25 of FIG. 2 (SEQ ID NO:2). Therefore, mature forms of the PRO19598 polypeptide which are encompassed by the present invention include those comprising amino acids X to 262 of FIG. 2 (SEQ ID NO:2), wherein X is any amino acid from 16 to 25 of FIG. 2 (SEQ ID NO:2) and variants thereof as described below. Isolated nucleic acid molecules encoding these polypeptides are also contemplated.

Another embodiment is directed to fragments of a PRO19598 polypeptide coding sequence that may find use as, for example, hybridization probes or for encoding fragments of a PRO19598 polypeptide that may optionally encode a polypeptide comprising a binding site for an anti-PRO19598 antibody. Such nucleic acid fragments are usually at least about 20 nucleotides in length, alternatively at least about 30 nucleotides in length, alternatively at least about 40 nucleotides in length, alternatively at least about 50 nucleotides in length, alternatively at least about 60 nucleotides in length, alternatively at least about 70 nucleotides in length, alternatively at least about 80 nucleotides in length, alternatively at least about 90 nucleotides in length, alternatively at least about 100 nucleotides in length, alternatively at least about 110 nucleotides in length, alternatively at least about 120 nucleotides in length, alternatively at least about 130 nucleotides in length, alternatively at least about 140 nucleotides in length, alternatively at least about 150 nucleotides in length, alternatively at least about 160 nucleotides in length, alternatively at least about 170 nucleotides in length, alternatively at least about 180 nucleotides in length, alternatively at least about 190 nucleotides in length, alternatively at least about 200 nucleotides in length, alternatively at least about 250 nucleotides in length, alternatively at least about 300 nucleotides in length, alternatively at least about 350 nucleotides in length, alternatively at least about 400 nucleotides in length, alternatively at least about 450 nucleotides in length, alternatively at least about 500 nucleotides in length, alternatively at least about 600 nucleotides in length, alternatively at least about 700 nucleotides in length, alternatively at least about 800 nucleotides in length, alternatively at least about 900 nucleotides in length and alternatively at least about 1000 nucleotides in length, wherein in this context the term "about" means the referenced nucleotide sequence length plus or minus 10% of that referenced length. In a preferred embodiment, the nucleotide sequence fragment is derived from any coding region of the nucleotide sequence shown in FIG. 1 (SEQ ID NO:1). It is noted that novel fragments of a PRO19598 polypeptide-encoding nucleotide sequence may be determined in a routine manner by aligning the PRO19598 polypeptide-encoding nucleotide sequence with other known nucleotide sequences using any of a number of well known sequence alignment programs and determining which PRO19598 polypeptide-encoding nucleotide sequence fragment(s) are novel. All of such PRO19598 polypeptide-encoding nucleotide sequences are contemplated herein and can be determined without undue experimentation. Also contemplated are the PRO19598 polypeptide fragments encoded by these nucleotide molecule fragments, preferably those PRO19598 polypeptide fragments that comprise a binding site for an anti-PRO19598 antibody.

In an other embodiment, the invention provides a vector comprising a nucleotide sequence encoding PRO19598 or

its variants. The vector may comprise any of the isolated nucleic acid molecules hereinabove identified.

A host cell comprising such a vector is also provided. By way of example, the host cells may be CHO cells, *E. coli*, yeast, or Baculovirus-infected insect cells. A process for producing PRO19598 polypeptides is further provided and comprises culturing host cells under conditions suitable for expression of PRO19598 and recovering PRO19598 from the cell culture.

In another embodiment, the invention provides isolated PRO19598 polypeptide encoded by any of the isolated nucleic acid sequences hereinabove identified.

In a specific aspect, the invention provides isolated native sequence PRO19598 polypeptide, which in certain embodiments, includes an amino acid sequence comprising residues from 1 or 21 to about 262 of FIG. 2 (SEQ ID NO:2).

In another aspect, the invention concerns an isolated PRO19598 polypeptide, comprising an amino acid sequence having at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83% amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively at least about 90% amino acid sequence identity, alternatively at least about 91% amino acid sequence identity, alternatively at least about 92% amino acid sequence identity, alternatively at least about 93% amino acid sequence identity, alternatively at least about 94% amino acid sequence identity, alternatively at least about 95% amino acid sequence identity, alternatively at least about 96% amino acid sequence identity, alternatively at least about 97% amino acid sequence identity, alternatively at least about 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to the sequence of amino acid residues from 1 or 21 to about 262, inclusive, of FIG. 2 (SEQ ID NO:2).

In a further aspect, the invention concerns an isolated PRO19598 polypeptide comprising an amino acid sequence having at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83% amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively at least about 90% amino acid sequence identity, alternatively at least about 91% amino acid sequence identity, alternatively at least about 92% amino acid sequence identity, alternatively at least about 93% amino acid sequence identity, alternatively at least about 94% amino acid sequence identity, alternatively at least about 95% amino acid sequence identity, alternatively at least about 96% amino acid sequence identity, alternatively at least about 97% amino acid sequence identity, alternatively at least about 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence iden-

tity to an amino acid sequence encoded by the human protein cDNA deposited with the ATCC on Mar. 21, 2000 under ATCC Deposit No. PTA-1532 (DNA145887-2849). In a preferred embodiment, the isolated PRO19598 polypeptide comprises an amino acid sequence encoded by the human protein cDNA deposited with the ATCC on Mar. 21, 2000 under ATCC Deposit No. PTA-1532 (DNA145887-2849).

In a further aspect, the invention concerns an isolated PRO19598 polypeptide comprising an amino acid sequence scoring at least about 80% positives, alternatively at least about 81% positives, alternatively at least about 82% positives, alternatively at least about 83% positives, alternatively at least about 84% positives, alternatively at least about 85% positives, alternatively at least about 86% positives, alternatively at least about 87% positives, alternatively at least about 88% positives, alternatively at least about 89% positives, alternatively at least about 90% positives, alternatively at least about 91% positives, alternatively at least about 92% positives, alternatively at least about 93% positives, alternatively at least about 94% positives, alternatively at least about 95% positives, alternatively at least about 96% positives, alternatively at least about 97% positives, alternatively at least about 98% positives and alternatively at least about 99% positives when compared with the amino acid sequence of residues from 1 or 21 to about 262, inclusive, of FIG. 2 (SEQ ID NO:2).

In a specific aspect, the invention provides an isolated PRO19598 polypeptide without the N-terminal signal sequence and/or the initiating methionine and is encoded by a nucleotide sequence that encodes such an amino acid sequence as hereinbefore described. Processes for producing the same are also herein described, wherein those processes comprise culturing a host cell comprising a vector which comprises the appropriate encoding nucleic acid molecule under conditions suitable for expression of the PRO19598 polypeptide and recovering the PRO19598 polypeptide from the cell culture.

In yet another aspect, the invention concerns an isolated PRO19598 polypeptide, comprising the sequence of amino acid residues from 1 or 21 to about 262, inclusive, of FIG. 2 (SEQ ID NO:2), or a fragment thereof which is biologically active or sufficient to provide a binding site for an anti-PRO19598 antibody, wherein the identification of PRO19598 polypeptide fragments that possess biological activity or provide a binding site for an anti-PRO19598 antibody may be accomplished in a routine manner using techniques which are well known in the art. Preferably, the PRO19598 fragment retains a qualitative biological activity of a native PRO19598 polypeptide.

In a still further aspect, the invention provides a polypeptide produced by (i) hybridizing a test DNA molecule under stringent conditions with (a) a DNA molecule encoding a PRO19598 polypeptide having the sequence of amino acid residues from 1 or 21 to about 262, inclusive, of FIG. 2 (SEQ ID NO:2), or (b) the complement of the DNA molecule of (a), and if the test DNA molecule has at least about an 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about

90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity to (a) or (b), (ii) culturing a host cell comprising the test DNA molecule under conditions suitable for expression of the polypeptide, and (iii) recovering the polypeptide from the cell culture.

In another embodiment, the invention provides chimeric molecules comprising a PRO19598 polypeptide fused to a heterologous polypeptide or amino acid sequence, wherein the PRO19598 polypeptide may comprise any PRO19598 polypeptide, variant or fragment thereof as hereinbefore described. An example of such a chimeric molecule comprises a PRO19598 polypeptide fused to an epitope tag sequence or a Fc region of an immunoglobulin.

In another embodiment, the invention provides an antibody as defined below which specifically binds to a PRO19598 polypeptide as hereinbefore described. Optionally, the antibody is a monoclonal antibody, an antibody fragment or a single chain antibody.

In yet another embodiment, the invention concerns agonists and antagonists of a native PRO19598 polypeptide as defined below. In a particular embodiment, the agonist or antagonist is an anti-PRO19598 antibody or a small molecule.

In a further embodiment, the invention concerns a method of identifying agonists or antagonists to a PRO19598 polypeptide which comprise contacting the PRO19598 polypeptide with a candidate molecule and monitoring a biological activity mediated by said PRO19598 polypeptide. Preferably, the PRO19598 polypeptide is a native PRO19598 polypeptide.

In a still further embodiment, the invention concerns a composition of matter comprising a PRO19598 polypeptide, or an agonist or antagonist of a PRO19598 polypeptide as herein described, or an anti-PRO19598 antibody, in combination with a carrier. Optionally, the carrier is a pharmaceutically acceptable carrier.

Another embodiment of the present invention is directed to the use of a PRO19598 polypeptide, or an agonist or antagonist thereof as herein described, or an anti-PRO19598 antibody, for the preparation of a medicament useful in the treatment of a condition which is responsive to the PRO19598 polypeptide, an agonist or antagonist thereof or an anti-PRO19598 antibody.

In yet other embodiments, the invention provides oligonucleotide probes which may be useful for isolating genomic and cDNA nucleotide sequences, measuring or detecting expression of an associated gene or as antisense probes, wherein those probes may be derived from any of the above or below described nucleotide sequences. Preferred probe lengths are described above.

In yet other embodiments, the present invention is directed to methods of using the PRO19598 polypeptides of the present invention for a variety of uses based upon the functional biological assay data presented in the Examples below. In a particular embodiment, a PRO19598 polypeptide functions as a cytokine receptor that acts as a Decoy receptor and a natural antagonist of a cytokine in the IL-10 family

[designated herein as a PRO3301 polypeptide which is encoded by a nucleic acid molecule designated herein as DNA88002].

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows a nucleotide sequence (SEQ ID NO:1) of a native sequence PRO19598 cDNA, wherein SEQ ID NO:1 is a clone designated herein as "DNA145887-2849".

FIG. 2 shows the amino acid sequence (SEQ ID NO:2) derived from the coding sequence of SEQ ID NO:1 shown in FIG. 1.

FIG. 3 shows a nucleotide sequence (SEQ ID NO:6) of a native sequence PRO3301 cDNA, wherein SEQ ID NO:3 is a clone designated herein as "DNA88002".

FIG. 4 shows the amino acid sequence (SEQ ID NO:7) derived from the coding sequence of SEQ ID NO:6 shown in FIG. 3.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

I. Definitions

The terms "PRO polypeptide" and "PRO" as used herein and when immediately followed by a numerical designation refer to various polypeptides, wherein the complete designation (i.e., PRO/number) refers to specific polypeptide sequences as described herein. The terms "PRO/number polypeptide" and "PRO/number" wherein the term "number" is provided as an actual numerical designation as used herein encompass native sequence polypeptides and polypeptide variants (which are further defined herein). The PRO polypeptides described herein may be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant or synthetic methods. The term "PRO polypeptide" refers to each individual PRO/number polypeptide disclosed herein. All disclosures in this specification which refer to the "PRO polypeptide" refer to each of the polypeptides individually as well as jointly. For example, descriptions of the preparation of, purification of, derivation of, formation of antibodies to or against, administration of, compositions containing, treatment of a disease with, etc., pertain to each polypeptide of the invention individually. The term "PRO polypeptide" also includes variants of the PRO/number polypeptides disclosed herein.

A "native sequence PRO19598 polypeptide" comprises a polypeptide having the same amino acid sequence as the corresponding PRO19598 polypeptide derived from nature. Such native sequence PRO19598 polypeptides can be isolated from nature or can be produced by recombinant or synthetic means. The term "native sequence 19598 polypeptide" specifically encompasses naturally-occurring truncated or secreted forms of the specific PRO19598 polypeptide (e.g., an extracellular domain sequence), naturally-occurring variant forms (e.g., alternatively spliced forms) and naturally-occurring allelic variants of the polypeptide. In various embodiments of the invention, the native sequence PRO19598 polypeptides disclosed herein are mature or full-length native sequence polypeptides comprising the full-length amino acids sequences shown in the accompanying figures. Start and stop codons are shown in bold font and underlined in the figures. However, while the PRO19598 polypeptide disclosed in the accompanying figures are shown to begin with methionine residues designated herein as amino acid position 1 in the figures, it is conceivable and possible that other methionine residues located either upstream or downstream from the amino acid position

1 in the figures may be employed as the starting amino acid residue for the PRO19598 polypeptides.

The PRO19598 polypeptide “extracellular domain” or “ECD” refers to a form of the PRO19598 polypeptide which is essentially free of the transmembrane and cytoplasmic domains. Ordinarily, a PRO19598 polypeptide ECD will have less than 1% of such transmembrane and/or cytoplasmic domains and preferably, will have less than 0.5% of such domains. It will be understood that any transmembrane domains identified for the PRO19598 polypeptides of the present invention are identified pursuant to criteria routinely employed in the art for identifying that type of hydrophobic domain. The exact boundaries of a transmembrane domain may vary but most likely by no more than about 5 amino acids at either end of the domain as initially identified herein. Optionally, therefore, an extracellular domain of a PRO19598 polypeptide may contain from about 5 or fewer amino acids on either side of the transmembrane domain/extracellular domain boundary as identified in the Examples or specification and such polypeptides, with or without the associated signal peptide, and nucleic acid encoding them, are contemplated by the present invention.

The approximate location of the “signal peptides” of the various PRO19598 polypeptides disclosed herein are shown in the present specification and/or the accompanying figures. It is noted, however, that the C-terminal boundary of a signal peptide may vary, but most likely by no more than about 5 amino acids on either side of the signal peptide C-terminal boundary as initially identified herein, wherein the C-terminal boundary of the signal peptide may be identified pursuant to criteria routinely employed in the art for identifying that type of amino acid sequence element (e.g., Nielsen et al., *Prot. Eng.* 10:1–6 (1997) and von Heinje et al., *Nucl. Acids. Res.* 14:4683–4690 (1986)). Moreover, it is also recognized that, in some cases, cleavage of a signal sequence from a secreted polypeptide is not entirely uniform, resulting in more than one secreted species. These mature polypeptides, where the signal peptide is cleaved within no more than about 5 amino acids on either side of the C-terminal boundary of the signal peptide as identified herein, and the polynucleotides encoding them, are contemplated by the present invention.

“PRO19598 polypeptide variant” means an active PRO19598 polypeptide as defined above or below having at least about 80% amino acid sequence identity with a full-length native sequence PRO19598 polypeptide sequence as disclosed herein, a PRO19598 polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO19598 polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO19598 polypeptide sequence as disclosed herein. Such PRO19598 polypeptide variants include, for instance, PRO19598 polypeptides wherein one or more amino acid residues are added, or deleted, at the N- or C-terminus of the full-length native amino acid sequence. Ordinarily, a PRO19598 polypeptide variant will have at least about 80% amino acid sequence identity, alternatively at least about 81% amino acid sequence identity, alternatively at least about 82% amino acid sequence identity, alternatively at least about 83% amino acid sequence identity, alternatively at least about 84% amino acid sequence identity, alternatively at least about 85% amino acid sequence identity, alternatively at least about 86% amino acid sequence identity, alternatively at least about 87% amino acid sequence identity, alternatively at least about 88% amino acid sequence identity, alternatively at least about 89% amino acid sequence identity, alternatively

at least about 90% amino acid sequence identity, alternatively at least about 91% amino acid sequence identity, alternatively at least about 92% amino acid sequence identity, alternatively at least about 93% amino acid sequence identity, alternatively at least about 94% amino acid sequence identity, alternatively at least about 95% amino acid sequence identity, alternatively at least about 96% amino acid sequence identity, alternatively at least about 97% amino acid sequence identity, alternatively at least about 98% amino acid sequence identity and alternatively at least about 99% amino acid sequence identity to a full-length native sequence PRO19598 polypeptide sequence as disclosed herein, a PRO19598 polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO19598 polypeptide, with or without the signal peptide, as disclosed herein or any other specifically defined fragment of a full-length PRO19598 polypeptide sequence as disclosed herein. Ordinarily, PRO19598 variant polypeptides are at least about 10 amino acids in length, alternatively at least about 20 amino acids in length, alternatively at least about 30 amino acids in length, alternatively at least about 40 amino acids in length, alternatively at least about 50 amino acids in length, alternatively at least about 60 amino acids in length, alternatively at least about 70 amino acids in length, alternatively at least about 80 amino acids in length, alternatively at least about 90 amino acids in length, alternatively at least about 100 amino acids in length, alternatively at least about 150 amino acids in length, alternatively at least about 200 amino acids in length, alternatively at least about 300 amino acids in length, or more.

“Percent (%) amino acid sequence identity” with respect to the PRO polypeptide sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in the specific PRO polypeptide sequence, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full length of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are generated using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1 below. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Table 1 below has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, Calif. or may be compiled from the source code provided in Table 1 below. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

In situations where ALIGN-2 is employed for amino acid sequence comparisons, the % amino acid sequence identity of a given amino acid sequence A to, with, or against a given

amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

100 times the fraction X/Y

where X is the number of amino acid residues scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A. As examples of % amino acid sequence identity calculations using this method, Tables 2 and 3 demonstrate how to calculate the % amino acid sequence identity of the amino acid sequence designated "Comparison Protein" to the amino acid sequence designated "PRO", wherein "PRO" represents the amino acid sequence of a hypothetical PRO polypeptide of interest, "Comparison Protein" represents the amino acid sequence of a polypeptide against which the "PRO" polypeptide of interest is being compared, and "X", "Y" and "Z" each represent different hypothetical amino acid residues.

Unless specifically stated otherwise, all % amino acid sequence identity values used herein are obtained as described in the immediately preceding paragraph using the ALIGN-2 computer program. However, % amino acid sequence identity values may also be obtained as described below by using the WU-BLAST-2 computer program (Altschul et al., *Methods in Enzymology* 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, i.e., the adjustable parameters, are set with the following values: overlap span=1, overlap fraction=0.125, word threshold (T)=11, and scoring matrix=BLOSUM62. When WU-BLAST-2 is employed, a % amino acid sequence identity value is determined by dividing (a) the number of matching identical amino acid residues between the amino acid sequence of the PRO polypeptide of interest having a sequence derived from the native PRO polypeptide and the comparison amino acid sequence of interest (i.e., the sequence against which the PRO polypeptide of interest is being compared which may be a PRO variant polypeptide) as determined by WU-BLAST-2 by (b) the total number of amino acid residues of the PRO polypeptide of interest. For example, in the statement "a polypeptide comprising an the amino acid sequence A which has or having at least 80% amino acid sequence identity to the amino acid sequence B", the amino acid sequence A is the comparison amino acid sequence of interest and the amino acid sequence B is the amino acid sequence of the PRO polypeptide of interest.

Percent amino acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul et al., *Nucleic Acids Res.* 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov> or otherwise obtained from the National Institute of Health, Bethesda, Md. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask=yes, strand=all, expected occurrences=10, minimum low complexity length=15/5, multi-pass e-value=0.01, constant for multi-pass=25, dropoff for final gapped alignment=25 and scoring matrix=BLOSUM62.

In situations where NCBI-BLAST2 is employed for amino acid sequence comparisons, the % amino acid

sequence identity of a given amino acid sequence A to, with, or against a given amino acid sequence B (which can alternatively be phrased as a given amino acid sequence A that has or comprises a certain % amino acid sequence identity to, with, or against a given amino acid sequence B) is calculated as follows:

100 times the fraction X/Y

where X is the number of amino acid residues scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of A and B, and where Y is the total number of amino acid residues in B. It will be appreciated that where the length of amino acid sequence A is not equal to the length of amino acid sequence B, the % amino acid sequence identity of A to B will not equal the % amino acid sequence identity of B to A.

"PRO19598 variant polynucleotide" or "PRO19598 variant nucleic acid sequence" means a nucleic acid molecule which encodes an active PRO19598 polypeptide as defined below and which has at least about 80% nucleic acid sequence identity with a nucleotide acid sequence encoding a full-length native sequence PRO19598 polypeptide sequence as disclosed herein, a full-length native sequence PRO19598 polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO19598 polypeptide, with or without the signal peptide, as disclosed herein or any other fragment of a full-length PRO19598 polypeptide sequence as disclosed herein. Ordinarily, a PRO19598 variant polynucleotide will have at least about 80% nucleic acid sequence identity, alternatively at least about 81% nucleic acid sequence identity, alternatively at least about 82% nucleic acid sequence identity, alternatively at least about 83% nucleic acid sequence identity, alternatively at least about 84% nucleic acid sequence identity, alternatively at least about 85% nucleic acid sequence identity, alternatively at least about 86% nucleic acid sequence identity, alternatively at least about 87% nucleic acid sequence identity, alternatively at least about 88% nucleic acid sequence identity, alternatively at least about 89% nucleic acid sequence identity, alternatively at least about 90% nucleic acid sequence identity, alternatively at least about 91% nucleic acid sequence identity, alternatively at least about 92% nucleic acid sequence identity, alternatively at least about 93% nucleic acid sequence identity, alternatively at least about 94% nucleic acid sequence identity, alternatively at least about 95% nucleic acid sequence identity, alternatively at least about 96% nucleic acid sequence identity, alternatively at least about 97% nucleic acid sequence identity, alternatively at least about 98% nucleic acid sequence identity and alternatively at least about 99% nucleic acid sequence identity with a nucleic acid sequence encoding a full-length native sequence PRO19598 polypeptide sequence as disclosed herein, a full-length native sequence PRO19598 polypeptide sequence lacking the signal peptide as disclosed herein, an extracellular domain of a PRO19598 polypeptide, with or without the signal sequence, as disclosed herein or any other fragment of a full-length PRO19598 polypeptide sequence as disclosed herein. Variants do not encompass the native nucleotide sequence.

Ordinarily, PRO19598 variant polynucleotides are at least about 30 nucleotides in length, alternatively at least about 60 nucleotides in length, alternatively at least about 90 nucleotides in length, alternatively at least about 120 nucleotides in length, alternatively at least about 150 nucleotides in length, alternatively at least about 180 nucleotides in length, alternatively at least about 210 nucleotides in length, alter-

natively at least about 240 nucleotides in length, alternatively at least about 270 nucleotides in length, alternatively at least about 300 nucleotides in length, alternatively at least about 450 nucleotides in length, alternatively at least about 600 nucleotides in length, alternatively at least about 900 nucleotides in length, or more.

"Percent (%) nucleic acid sequence identity" with respect to PRO-encoding nucleic acid sequences identified herein is defined as the percentage of nucleotides in a candidate sequence that are identical with the nucleotides in the PRO nucleic acid sequence of interest, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity. Alignment for purposes of determining percent nucleic acid sequence identity can be achieved in various ways that are within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN or Megalign (DNASTAR) software. For purposes herein, however, % nucleic acid sequence identity values are generated using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in Table 1 below. The ALIGN-2 sequence comparison computer program was authored by Genentech, Inc. and the source code shown in Table 1 below has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco, Calif. or may be compiled from the source code provided in Table 1 below. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.

In situations where ALIGN-2 is employed for nucleic acid sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

$$100 \text{ times the fraction } W/Z$$

where W is the number of nucleotides scored as identical matches by the sequence alignment program ALIGN-2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C. As examples of % nucleic acid sequence identity calculations, Tables 4 and 5, demonstrate how to calculate the % nucleic acid sequence identity of the nucleic acid sequence designated "Comparison DNA" to the nucleic acid sequence designated "PRO-DNA", wherein "PRO-DNA" represents a hypothetical PRO-encoding nucleic acid sequence of interest, "Comparison DNA" represents the nucleotide sequence of a nucleic acid molecule against which the "PRO-DNA" nucleic acid molecule of interest is being compared, and "N", "L" and "V" each represent different hypothetical nucleotides.

Unless specifically stated otherwise, all % nucleic acid sequence identity values used herein are obtained as described in the immediately preceding paragraph using the ALIGN-2 computer program. However, % nucleic acid sequence identity values may also be obtained as described

below by using the WU-BLAST-2 computer program (Altschul et al., *Methods in Enzymology* 266:460-480 (1996)). Most of the WU-BLAST-2 search parameters are set to the default values. Those not set to default values, i.e., the adjustable parameters, are set with the following values: overlap span=1, overlap fraction=0.125, word threshold (T)=11, and scoring matrix=BLOSUM62. When WU-BLAST-2 is employed, a % nucleic acid sequence identity value is determined by dividing (a) the number of matching identical nucleotides between the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid molecule of interest having a sequence derived from the native sequence PRO polypeptide-encoding nucleic acid and the comparison nucleic acid molecule of interest (i.e., the sequence against which the PRO polypeptide-encoding nucleic acid molecule of interest is being compared which may be a variant PRO polynucleotide) as determined by WU-BLAST-2 by (b) the total number of nucleotides of the PRO polypeptide-encoding nucleic acid molecule of interest. For example, in the statement "an isolated nucleic acid molecule comprising a nucleic acid sequence A which has or having at least 80% nucleic acid sequence identity to the nucleic acid sequence B", the nucleic acid sequence A is the comparison nucleic acid molecule of interest and the nucleic acid sequence B is the nucleic acid sequence of the PRO polypeptide-encoding nucleic acid molecule of interest.

Percent nucleic acid sequence identity may also be determined using the sequence comparison program NCBI-BLAST2 (Altschul et al., *Nucleic Acids Res.* 25:3389-3402 (1997)). The NCBI-BLAST2 sequence comparison program may be downloaded from <http://www.ncbi.nlm.nih.gov> or otherwise obtained from the National Institute of Health, Bethesda, Md. NCBI-BLAST2 uses several search parameters, wherein all of those search parameters are set to default values including, for example, unmask=yes, strand=all, expected occurrences=10, minimum low complexity length=15/5, multi-pass e-value =0.01, constant for multi-pass=25, dropoff for final gapped alignment=25 and scoring matrix=BLOSUM62.

In situations where NCBI-BLAST2 is employed for sequence comparisons, the % nucleic acid sequence identity of a given nucleic acid sequence C to, with, or against a given nucleic acid sequence D (which can alternatively be phrased as a given nucleic acid sequence C that has or comprises a certain % nucleic acid sequence identity to, with, or against a given nucleic acid sequence D) is calculated as follows:

$$100 \text{ times the fraction } W/Z$$

where W is the number of nucleotides scored as identical matches by the sequence alignment program NCBI-BLAST2 in that program's alignment of C and D, and where Z is the total number of nucleotides in D. It will be appreciated that where the length of nucleic acid sequence C is not equal to the length of nucleic acid sequence D, the % nucleic acid sequence identity of C to D will not equal the % nucleic acid sequence identity of D to C.

In other embodiments, PRO19598 variant polynucleotides are nucleic acid molecules that encode an active PRO19598 polypeptide and which are capable of hybridizing, preferably under stringent hybridization and wash conditions, to nucleotide sequences encoding a full-length PRO19598 polypeptide as disclosed herein. PRO19598 variant polypeptides may be those that are encoded by a PRO19598 variant polynucleotide.

"Isolated," when used to describe the various polypeptides disclosed herein, means polypeptide that has been

identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the polypeptide will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Isolated polypeptide includes polypeptide in situ within recombinant cells, since at least one component of the PRO19598 polypeptide natural environment will not be present. Ordinarily, however, isolated polypeptide will be prepared by at least one purification step.

An "isolated" PRO19598 polypeptide-encoding nucleic acid or other polypeptide-encoding nucleic acid is a nucleic acid molecule that is identified and separated from at least one contaminant nucleic acid molecule with which it is ordinarily associated in the natural source of the polypeptide-encoding nucleic acid. An isolated polypeptide-encoding nucleic acid molecule is other than in the form or setting in which it is found in nature. Isolated polypeptide-encoding nucleic acid molecules therefore are distinguished from the specific polypeptide-encoding nucleic acid molecule as it exists in natural cells. However, an isolated polypeptide-encoding nucleic acid molecule includes polypeptide-encoding nucleic acid molecules contained in cells that ordinarily express the polypeptide where, for example, the nucleic acid molecule is in a chromosomal location different from that of natural cells.

The term "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

The term "antibody" is used in the broadest sense and specifically covers, for example, single anti-PRO19598 monoclonal antibodies (including agonist, antagonist, and neutralizing antibodies), anti-PRO19598 antibody compositions with polypeptidic specificity, single chain anti-PRO19598 antibodies, and fragments of anti-PRO19598 antibodies (see below). The term "monoclonal antibody" as used herein refers to an antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that may be present in minor amounts.

"Stringency" of hybridization reactions is readily determinable by one of ordinary skill in the art, and generally is an empirical calculation dependent upon probe length, washing temperature, and salt concentration. In general, longer probes require higher temperatures for proper annealing, while shorter probes need lower temperatures. Hybridization generally depends on the ability of denatured DNA to reanneal when complementary strands are present in an environment below their melting temperature. The higher the degree of desired homology between the probe and hybridizable sequence, the higher the relative temperature which can be used. As a result, it follows that higher relative temperatures would tend to make the reaction conditions more stringent, while lower temperatures less so. For additional details and explanation of stringency of hybridization reactions, see Ausubel et al., *Current Protocols in Molecular Biology*, Wiley Interscience Publishers, (1995).

"Stringent conditions" or "high stringency conditions", as defined herein, may be identified by those that: (1) employ low ionic strength and high temperature for washing, for example 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate at 50° C.; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (v/v) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42° C.; or (3) employ 50% formamide, 5×SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5×Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42° C., with washes at 42° C. in 0.2×SSC (sodium chloride/sodium citrate) and 50% formamide at 55° C., followed by a high-stringency wash consisting of 0.1×SSC containing EDTA at 55° C.

"Moderately stringent conditions" may be identified as described by Sambrook et al., *Molecular Cloning: A Laboratory Manual*, New York: Cold Spring Harbor Press, 1989, and include the use of washing solution and hybridization conditions (e.g., temperature, ionic strength and % SDS) less stringent than those described above. An example of moderately stringent conditions is overnight incubation at 37° C. in a solution comprising: 20% formamide, 5×SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5×Denhardt's solution, 10% dextran sulfate, and 20 mg/ml denatured sheared salmon sperm DNA, followed by washing the filters in 1×SSC at about 37–50° C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc. as necessary to accommodate factors such as probe length and the like.

The term "epitope tagged" when used herein refers to a chimeric polypeptide comprising a PRO19598 polypeptide fused to a "tag polypeptide". The tag polypeptide has enough residues to provide an epitope against which an antibody can be made, yet is short enough such that it does not interfere with activity of the polypeptide to which it is fused. The tag polypeptide preferably also is fairly unique so that the antibody does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have at least six amino acid residues and usually between about 8 and 50 amino acid residues (preferably, between about 10 and 20 amino acid residues).

As used herein, the term "immunoadhesin" designates antibody-like molecules which combine the binding specificity of a heterologous protein (an "adhesin") with the effector functions of immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of an

amino acid sequence with the desired binding specificity which is other than the antigen recognition and binding site of an antibody (i.e., is "heterologous"), and an immunoglobulin constant domain sequence. The adhesin part of an immunoadhesin molecule typically is a contiguous amino acid sequence comprising at least the binding site of a receptor or a ligand. The immunoglobulin constant domain sequence in the immunoadhesin may be obtained from any immunoglobulin, such as IgG-1, IgG-2, IgG-3, or IgG-4 subtypes, IgA (including IgA-1 and IgA-2), IgE, IgD or IgM.

"Active" or "activity" for the purposes herein refers to form(s) of a PRO19598 polypeptide which retain a biological and/or an immunological activity of native or naturally-occurring PRO19598, wherein "biological" activity refers to a biological function (either inhibitory or stimulatory) caused by a native or naturally-occurring PRO19598 other than the ability to induce the production of an antibody against an antigenic epitope possessed by a native or naturally-occurring PRO19598 and an "immunological" activity refers to the ability to induce the production of an antibody against an antigenic epitope possessed by a native or naturally-occurring PRO19598.

The term "antagonist" is used in the broadest sense, and includes any molecule that partially or fully blocks, inhibits, or neutralizes a biological activity of a native PRO19598 polypeptide disclosed herein. In a similar manner, the term "agonist" is used in the broadest sense and includes any molecule that mimics a biological activity of a native PRO19598 polypeptide disclosed herein. Suitable agonist or antagonist molecules specifically include agonist or antagonist antibodies or antibody fragments, fragments or amino acid sequence variants of native PRO19598 polypeptides, peptides, antisense oligonucleotides, small organic molecules, etc. Methods for identifying agonists or antagonists of a PRO19598 polypeptide may comprise contacting a PRO19598 polypeptide with a candidate agonist or antagonist molecule and measuring a detectable change in one or more biological activities normally associated with the PRO19598 polypeptide.

"Treatment" refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) the targeted pathologic condition or disorder. Those in need of treatment include those already with the disorder as well as those prone to have the disorder or those in whom the disorder is to be prevented.

"Chronic" administration refers to administration of the agent(s) in a continuous mode as opposed to an acute mode, so as to maintain the initial therapeutic effect (activity) for an extended period of time. "Intermittent" administration is treatment that is not consecutively done without interruption, but rather is cyclic in nature.

"Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, cats, cattle, horses, sheep, pigs, goats, rabbits, etc. Preferably, the mammal is human.

Administration "in combination with" one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order.

"Carriers" as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers which are non-toxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically acceptable carrier is an aqueous pH buffered solution. Examples of physiologically acceptable carriers include buffers such as phosphate, citrate, and other organic acids;

antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN™, polyethylene glycol (PEG), and PLURONICS™.

"Antibody fragments" comprise a portion of an intact antibody, preferably the antigen binding or variable region of the intact antibody. Examples of antibody fragments include Fab, Fab', F(ab')₂, and Fv fragments; diabodies; linear antibodies (Zapata et al., *Protein Eng.* 8(10): 1057-1062 [1995]); single-chain antibody molecules; and multispecific antibodies formed from antibody fragments.

Papain digestion of antibodies produces two identical antigen-binding fragments, called "Fab" fragments, each with a single antigen-binding site, and a residual "Fc" fragment, a designation reflecting the ability to crystallize readily. Pepsin treatment yields an F(ab')₂ fragment that has two antigen-combining sites and is still capable of cross-linking antigen.

"Fv" is the minimum antibody fragment which contains a complete antigen-recognition and -binding site. This region consists of a dimer of one heavy- and one light-chain variable domain in tight, non-covalent association. It is in this configuration that the three CDRs of each variable domain interact to define an antigen-binding site on the surface of the V_H-V_L dimer. Collectively, the six CDRs confer antigen-binding specificity to the antibody. However, even a single variable domain (or half of an Fv comprising only three CDRs specific for an antigen) has the ability to recognize and bind antigen, although at a lower affinity than the entire binding site.

The Fab fragment also contains the constant domain of the light chain and the first constant domain (CH1) of the heavy chain. Fab fragments differ from Fab' fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH1 domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free thiol group. F(ab')₂ antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

The "light chains" of antibodies (immunoglobulins) from any vertebrate species can be assigned to one of two clearly distinct types, called kappa and lambda, based on the amino acid sequences of their constant domains.

Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG, and IgM, and several of these may be further divided into subclasses (isotypes), e.g., IgG1, IgG2, IgG3, IgG4, IgA, and IgA2.

"Single-chain Fv" or "sFv" antibody fragments comprise the V_H and V_L domains of antibody, wherein these domains are present in a single polypeptide chain. Preferably, the Fv polypeptide further comprises a polypeptide linker between the V_H and V_L domains which enables the sFv to form the desired structure for antigen binding. For a review of sFv, see Pluckthun in *The Pharmacology of Monoclonal Antibodies*, vol. 113, Rosenberg and Moore eds., Springer-Verlag, New York, pp. 269-315 (1994).

The term "diabodies" refers to small antibody fragments with two antigen-binding sites, which fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) in the same polypeptide chain (V_H-V_L). By using a linker that is too short to allow pairing between the two domains on the same chain, the domains are forced to pair with the complementary domains of another chain and create two antigen-binding sites. Diabodies are described more fully in, for example, EP 404,097; WO 93/11161; and Hollinger et al., *Proc. Natl. Acad. Sci. USA*, 90:6444-6448 (1993).

An "isolated" antibody is one which has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials which would interfere with diagnostic or therapeutic uses for the antibody, and may include enzymes, hormones, and other proteinaceous or nonproteinaceous solutes. In preferred embodiments, the antibody will be purified (1) to greater than 95% by weight of antibody as determined by the Lowry method, and most preferably more than 99% by weight, (2) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (3) to homogeneity by SDS-PAGE under reducing or nonreducing conditions using Coomassie blue or, preferably, silver stain. Isolated antibody includes the antibody in situ within recombinant cells since at least one component of the antibody's natural environment will not be present. Ordinarily, however, isolated antibody will be prepared by at least one purification step.

An antibody that "specifically binds to" or is "specific for" a particular polypeptide or an epitope on a particular polypeptide is one that binds to that particular polypeptide or

epitope on a particular polypeptide without substantially binding to any other polypeptide or polypeptide epitope.

The word "label" when used herein refers to a detectable compound or composition which is conjugated directly or indirectly to the antibody so as to generate a "labeled" antibody. The label may be detectable by itself (e.g. radioisotope labels or fluorescent labels) or, in the case of an enzymatic label, may catalyze chemical alteration of a substrate compound or composition which is detectable.

By "solid phase" is meant a non-aqueous matrix to which the antibody of the present invention can adhere. Examples of solid phases encompassed herein include those formed partially or entirely of glass (e.g., controlled pore glass), polysaccharides (e.g., agarose), polyacrylamides, polystyrene, polyvinyl alcohol and silicones. In certain embodiments, depending on the context, the solid phase can comprise the well of an assay plate; in others it is a purification column (e.g., an affinity chromatography column). This term also includes a discontinuous solid phase of discrete particles, such as those described in U.S. Pat. No. 4,275,149.

A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant which is useful for delivery of a drug (such as a PRO19598 polypeptide or antibody thereto) to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

A "small molecule" is defined herein to have a molecular weight below about 500 Daltons.

An "effective amount" of a polypeptide disclosed herein or an agonist or antagonist thereof is an amount sufficient to carry out a specifically stated purpose. An "effective amount" may be determined empirically and in a routine manner, in relation to the stated purpose.

TABLE 1

/*	
.	
* C-C increased from 12 to 15	
* Z is average of EQ	
* B is average of ND	
* match with stop is _M; stop-stop = 0; J (joker) match = 0	
*/	
#define _M	-8 /* value of a match with a stop */
int	_day[26][26] = {
/*	
A B C D E F G H I J K L M N O P Q R S T U V W X Y Z	
/* A */	{2, 0, -2, 0, 0, -4, 1, -1, -1, 0, -1, -2, -1, 0, _M, 1, 0, -2, 1, 1, 0, 0, -6, 0, -3, 0},
/* B */	{0, 3, -4, 3, 2, -5, 0, 1, -2, 0, 0, -3, -2, 2, _M, -1, 1, 0, 0, 0, -2, -5, 0, -3, 1},
/* C */	{-2, -4, 15, -5, -5, -4, -3, -3, -2, 0, -5, -6, -5, -4, _M, -3, -5, -4, 0, -2, 0, -2, -8, 0, 0, -5},
/* D */	{0, 3, -5, 4, 3, -6, 1, 1, -2, 0, 0, -4, -3, 2, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 2},
/* E */	{0, 2, -5, 3, 4, -5, 0, 1, -2, 0, 0, -3, -2, 1, _M, -1, 2, -1, 0, 0, 0, -2, -7, 0, -4, 3},
/* F */	{-4, -5, -4, -6, -5, 9, -5, -2, 1, 0, -5, 2, 0, -4, _M, -5, -5, -4, -3, -3, 0, -1, 0, 0, 7, -5},
/* G */	{1, 0, -3, 1, 0, -5, 5, -2, -3, 0, -2, -4, -3, 0, _M, -1, -1, -3, 1, 0, 0, -1, -7, 0, -5, 0},
/* H */	{-1, 1, -3, 1, 1, -2, -2, 6, -2, 0, 0, -2, -2, 2, _M, 0, 3, 2, -1, -1, 0, -2, -3, 0, 0, 2},
/* I */	{-1, -2, -2, -2, 1, -3, -2, 5, 0, -2, 2, 2, -2, _M, -2, -2, -2, -1, 0, 0, 4, -5, 0, -1, -2},
/* J */	{0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* K */	{-1, 0, -5, 0, 0, -5, -2, 0, -2, 0, 5, -3, 0, 1, _M, -1, 1, 3, 0, 0, 0, -2, -3, 0, -4, 0},
/* L */	{-2, -3, -6, -4, -3, 2, -4, -2, 2, 0, -3, 6, 4, -3, _M, -3, -2, -3, -3, -1, 0, 2, -2, 0, -1, -2},
/* M */	{-1, -2, -5, -3, -2, 0, -3, -2, 2, 0, 0, 4, 6, -2, _M, -2, -1, 0, -2, -1, 0, 2, -4, 0, -2, -1},
/* N */	{0, 2, -4, 2, 1, -4, 0, 2, -2, 0, 1, -3, -2, 2, _M, -1, 1, 0, 1, 0, 0, -2, -4, 0, -2, 1},
/* O */	{_M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, _M, 0, _M, _M, _M, _M, _M, _M, _M, _M, _M},
/* P */	{1, -1, -3, -1, -1, -5, -1, 0, -2, 0, -1, -3, -2, -1, _M, 6, 0, 0, 1, 0, 0, -1, -6, 0, -5, 0},
/* Q */	{0, 1, -5, 2, 2, -5, -1, 3, -2, 0, 1, -2, -1, 1, _M, 0, 4, 1, -1, -1, 0, -2, -5, 0, -4, 3},
/* R */	{-2, 0, -4, -1, -1, -4, -3, 2, -2, 0, 3, -3, 0, 0, _M, 0, 1, 6, 0, -1, 0, -2, 2, 0, -4, 0},
/* S */	{1, 0, 0, 0, 0, -3, 1, -1, -1, 0, 0, -3, -2, 1, _M, 1, -1, 0, 2, 1, 0, -1, -2, 0, -3, 0},
/* T */	{1, 0, -2, 0, 0, -3, 0, -1, 0, 0, 0, -1, -1, 0, _M, 0, -1, -1, 1, 3, 0, 0, -5, 0, -3, 0},
/* U */	{0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* V */	{0, -2, -2, -2, -2, -1, -1, -2, 4, 0, -2, 2, 2, -2, _M, -1, -2, -2, -1, 0, 0, 4, -6, 0, -2, -2},
/* W */	{-6, -5, -8, -7, -7, 0, -7, -3, -5, 0, -3, -2, -4, -4, _M, -6, -5, 2, -2, -5, 0, -6, 17, 0, 0, -6},
/* X */	{0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0, _M, 0, 0, 0, 0, 0, 0, 0, 0, 0, 0},
/* Y */	{-3, -3, 0, -4, -4, 7, -5, 0, -1, 0, -4, -1, -2, -2, _M, -5, -4, -4, -3, -3, 0, -2, 0, 0, 10, -4},
/* Z */	{0, 1, -5, 2, 3, -5, 0, 2, -2, 0, 0, -2, -1, 1, _M, 0, 3, 0, 0, 0, -2, -6, 0, -4, 4},
};	

TABLE 1-continued

```

/*
*/
#include <stdio.h>
#include <ctype.h>
#define MAXJMP      16 /* max jumps in a diag */
#define MAXGAP      24 /* don't continue to penalize gaps larger than this */
#define JMPS        1024 /* max jmps in an path */
#define MX          4 /* save if there's at least MX-1 bases since last jmp */
#define DMAT        3 /* value of matching bases */
#define DMIS        0 /* penalty for mismatched bases */
#define DINS0       8 /* penalty for a gap */
#define DINS1       1 /* penalty per base */
#define PINS0       8 /* penalty for a gap */
#define PINS1       4 /* penalty per residue */
struct jmp {
    short      n[MAXJMP]; /* size of jmp (neg for dely) */
    unsigned short x[MAXJMP]; /* base no. of jmp in seq x */
    /* limits seq to 2 16 -1 */
};
struct diag {
    int      score; /* score at last jmp */
    long     offset; /* offset of prev block */
    short    ijmp; /* current jmp index */
    struct jmp jp; /* list of jmps */
};
struct path {
    int      spc; /* number of leading spaces */
    short    n[JMPS]; /* size of jmp (gap) */
    int      x[JMPS]; /* loc of jmp (last elem before gap) */
};
char      *ofile; /* output file name */
char      *namex[2]; /* seq names: getseqs() */
char      *prog; /* prog name for err msgs */
char      *seqx[2]; /* seqs: getseqs() */
int      dmax; /* best diag: nw() */
int      dmax0; /* final diag */
int      dna; /* set if dna: main() */
int      endgaps; /* set if penalizing end gaps */
int      gapx, gapy; /* total gaps in seqs */
int      len0, len1; /* seq lens */
int      ngapx, ngapy; /* total size of gaps */
int      smax; /* max score: nw() */
int      *xbm; /* bitmap for matching */
long     offset; /* current offset in jmp file */
struct diag *dx; /* holds diagonals */
struct path pp[2]; /* holds path for seqs */
char      *calloc(), *malloc(), *index(), *strcpy();
char      *getseq(), *g_calloc();
/* Needleman-Wunsch alignment program
*
* usage: progs file1 file2
* where file1 and file2 are two dna or two protein sequences.
* The sequences can be in upper- or lower-case an may contain ambiguity
* Any lines beginning with ';', '>' or '<' are ignored
* Max file length is 65535 (limited by unsigned short x in the jmp struct)
* A sequence with 1/4 or more of its elements ACGTU is assumed to be DNA
* Output is in the file "align.out"
*
* The program may create a tmp file in /tmp to hold info about traceback.
* Original version developed under BSD 4.3 on a vax 8650
*/
#include "nw.h"
#include "day.h"
static __dbval[26] = {
    1,14,2,13,0,0,4,11,0,0,12,0,3,15,0,0,0,5,6,8,8,7,9,0,10,0
};
static __pbval[26] = {
    1, 2[(1<<'D'-'A')](1<<'N'-'A')), 4, 8, 16, 32, 64,
    128, 256, 0xFFFFFFF, 1<<10, 1<<11, 1<<12, 1<<13, 1<<14,
    1<<15, 1<<16, 1<<17, 1<<18, 1<<19, 1<<20, 1<<21, 1<<22,
    1<<23, 1<<24, 1<<25[(1<<'E'-'A')](1<<'Q'-'A'))
};
main(ac, av)
int      ac;
char      *av[];
{
    prog = av[0];
    if(ac != 3) {
        fprintf(stderr, "usage: %s file1 file2\n", prog);
    }
}

```

main

TABLE 1-continued

```

fprintf(stderr, "where file1 and file2 are two dna or two protein sequences.\n");
fprintf(stderr, "The sequences can be in upper- or lower-case\n");
fprintf(stderr, "Any lines beginning with ';' or '<' are ignored\n");
fprintf(stderr, "Output is in the file \"align.out\"\n");
exit(1);
}
namex[0] = av[1];
namex[1] = av[2];
seqx[0] = getseq(namex[0], &len0);
seqx[1] = getseq(namex[1], &len1);
xbm = (dna)? _dbval : _pbval;
endgaps = 0; /* 1 to penalize endgaps */
ofile = "align.out"; /* output file */
nw(); /* fill in the matrix, get the possible jumps */
readjumps(); /* get the actual jumps */
print(); /* print stats, alignment */
cleanup(); /* unlink any tmp files */
}
/* do the alignment, return best score: main()
 * dna: values in Fitch and Smith, PNAS, 80, 1382-1386, 1983
 * pro: PAM 250 values
 * When scores are equal, we prefer mismatches to any gap, prefer
 * a new gap to extending an ongoing gap, and prefer a gap in seqx
 * to a gap in seq y.
 */
nw()
{
    char *px, *py; /* seqs and ptrs */
    int *ndely, *dely; /* keep track of dely */
    int ndelx, delx; /* keep track of delx */
    int *tmp; /* for swapping row0, row1 */
    int mis; /* score for each type */
    int ins0, ins1; /* insertion penalties */
    register id; /* diagonal index */
    register ij; /* jmp index */
    register *col0, *col1; /* score for curr, last row */
    register xx, yy; /* index into seqs */
    dx = (struct diag *)g_calloc("to get diag", len0+len1+1, sizeof(struct diag));
    ndely = (int *)g_calloc("to get ndely", len1+1, sizeof(int));
    dely = (int *)g_calloc("to get dely", len1+1, sizeof(int));
    col0 = (int *)g_calloc("to get col0", len1+1, sizeof(int));
    col1 = (int *)g_calloc("to get col1", len1+1, sizeof(int));
    ins0 = (dna)? DINS0 : PINS0;
    ins1 = (dna)? DINS1 : PINS1;
    smax = -10000;
    if (endgaps) {
        for (col0[0] = dely[0] = -ins0, yy = 1; yy <= len1; yy++) {
            col0[yy] = dely[yy] = col0[yy-1] - ins1;
            ndely[yy] = yy;
        }
        col0[0] = 0; /* Waterman Bull Math Biol 84 */
    }
    else
        for (yy = 1; yy <= len1; yy++)
            dely[yy] = -ins0;
    /* fill in match matrix
     */
    for (px = seqx[0], xx = 1; xx <= len0; px++, xx++) {
        /* initialize first entry in col
         */
        if (endgaps) {
            if (xx == 1)
                col1[0] = delx = -(ins0+ins1);
            else
                col1[0] = delx = col0[0]-ins1;
            ndelx = xx;
        }
        else {
            col1[0] = 0;
            delx = -ins0;
            ndelx = 0;
        }
    }

    for (py = seqx[1], yy = 1; yy <= len1; py++, yy++) {
        mis = col0[yy-1];
        if (dna)
            mis += (xbm["px-'A']&xbm["py-'A']? DMAT : DMIS;
        else
            mis += _day["px-'A']["py-'A'];
    }
}

```

...nw

TABLE 1-continued

```

/* update penalty for del in x seq;
 * favor new del over ongoing del
 * ignore MAXGAP if weighting endgaps
 */
if (endgaps || ndely[yy] < MAXGAP) {
    if (col0[yy] - ins0 >= dely[yy]) {
        dely[yy] = col0[yy] - (ins0+ins1);
        ndely[yy] = 1;
    } else {
        dely[yy] -= ins1;
        ndely[yy]++;
    }
} else {
    if (col0[yy] - (ins0+ins1) >= dely[yy]) {
        dely[yy] = col0[yy] - (ins0+ins1);
        ndely[yy] = 1;
    } else
        ndely[yy]++;
}
/* update penalty for del in y seq;
 * favor new del over ongoing del
 */
if (endgaps || ndelx < MAXGAP) {
    if (col1[yy-1] - ins0 >= delx) {
        delx = col1[yy-1] - (ins0+ins1);
        ndelx = 1;
    } else {
        delx -= ins1;
        ndelx++;
    }
} else {
    if (col1[yy-1] - (ins0+ins1) >= delx) {
        delx = col1[yy-1] - (ins0+ins1);
        ndelx = 1;
    } else
        ndelx++;
}
/* pick the maximum score; we're favoring
 * mis over any del and delx over dely
 */

id = xx - yy + len1 - 1;
if (mis >= delx && mis >= dely[yy])
    col1[yy] = mis;
else if (delx >= dely[yy]) {
    col1[yy] = delx;
    ij = dx[id].ijmp;
    if (dx[id].jp.n[0] && (!dna || (ndelx >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
        dx[id].ijmp++;
        if (++ij >= MAXJMP) {
            writejmps(id);
            ij = dx[id].ijmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
    }
    dx[id].jp.n[ij] = ndelx;
    dx[id].jp.x[ij] = xx;
    dx[id].score = delx;
}
else {
    col1[yy] = dely[yy];
    ij = dx[id].ijmp;
    if (dx[id].jp.n[0] && (!dna || (ndely[yy] >= MAXJMP
    && xx > dx[id].jp.x[ij]+MX) || mis > dx[id].score+DINS0)) {
        dx[id].ijmp++;
        if (++ij >= MAXJMP) {
            writejmps(id);
            ij = dx[id].ijmp = 0;
            dx[id].offset = offset;
            offset += sizeof(struct jmp) + sizeof(offset);
        }
    }
    dx[id].jp.n[ij] = ndely[yy];
    dx[id].jp.x[ij] = xx;
    dx[id].score = dely[yy];
}
if (xx == len0 && yy < len1) {

```

TABLE 1-continued

```

        /* last col
        */
        if (endgaps)
            col1[yy] -= ins0+ins1*(len1-yy);
        if(col1[yy] > smax) {
            smax = col1[yy];
            dmax = id;
        }
    }
}
if (endgaps && xx < len0)
    col1[yy-1] -= ins0+ins1*(len0-xx);
if (col1[yy-1] > smax) {
    smax = col1[yy-1];
    dmax = id;
}
tmp = col0; col0 = col1; col1 = tmp;
}
(void) free((char *)ndely);
(void) free((char *)dely);
(void) free((char *)col0);
(void) free((char *)col1);
}
/*
*
* print() -- only routine visible outside this module
*
* static:
* getmat() -- trace back best path, count matches: print()
* pr_align() -- print alignment of described in array p[]: print()
* dumpblock() -- dump a block of lines with numbers, stars: pr_align()
* nums() -- put out a number line: dumpblock()
* putline() -- put out a line (name, [num], seq, [num]): dumpblock()
* stars() -- put a line of stars: dumpblock()
* stripname() -- strip any path and prefix from a seqname
*/
#include "nw.h"
#define SPC 3
#define P__LINE 256 /* maximum output line */
#define P__SPC 3 /* space between name or num and seq */
extern _day[26][26];
int olen; /* set output line length */
FILE *fx; /* output file */
print()
{
    int lx, ly, firstgap, lastgap; /* overlap */
    if ((fx = fopen(ofile, "w")) == 0) {
        fprintf(stderr, "%s: can't write %s\n", prog, ofile);
        cleanup(1);
    }
    fprintf(fx, "<first sequence: %s (length = %d)\n", namex[0], len0);
    fprintf(fx, "<second sequence: %s (length = %d)\n", namex[1], len1);
    olen = 60;
    lx = len0;
    ly = len1;
    firstgap = lastgap = 0;
    if (dmax < len1 - 1) { /* leading gap in x */
        pp[0].spc = firstgap = len1 - dmax - 1;
        ly -= pp[0].spc;
    }
    else if (dmax > len1 - 1) { /* leading gap in y */
        pp[1].spc = firstgap = dmax - (len1 - 1);
        lx -= pp[1].spc;
    }
    if (dmax0 < len0 - 1) { /* trailing gap in x */
        lastgap = len0 - dmax0 - 1;
        lx -= lastgap;
    }
    else if (dmax0 > len0 - 1) { /* trailing gap in y */
        lastgap = dmax0 - (len0 - 1);
        ly -= lastgap;
    }
    getmat(lx, ly, firstgap, lastgap);
    pr_align();
}
/*
* trace back the best path, count matches
*/
static

```

TABLE 1-continued

```

getmat(lx, ly, firstgap, lastgap)                                     getmat
int      lx, ly;                                                    /* "core" (minus endgaps) */
int      firstgap, lastgap;                                          /* leading trailing overlap */
{
    int      nm, i0, i1, siz0, siz1;
    char      outx[32];
    double    pct;
    register  n0, n1;
    register char *p0, *p1;
    /* get total matches, score
    */
    i0 = i1 = siz0 = siz1 = 0;
    p0 = seqx[0] + pp[1].spc;
    p1 = seqx[1] + pp[0].spc;
    n0 = pp[1].spc + 1;
    n1 = pp[0].spc + 1;
    nm = 0;
    while ( *p0 && *p1 ) {
        if (siz0) {
            p1++;
            n1++;
            siz0--;
        }
        else if (siz1) {
            p0++;
            n0++;
            siz1--;
        }
        else {
            if (xbm[*p0-'A'] & xbm[*p1-'A'])
                nm++;
            if (n0++ == pp[0].x[i0])
                siz0 = pp[0].n[i0++];
            if (n1++ == pp[1].x[i1])
                siz1 = pp[1].n[i1++];
            p0++;
            p1++;
        }
    }
    /* pct homology:
    * if penalizing endgaps, base is the shorter seq
    * else, knock off overhangs and take shorter core
    */
    if (endgaps)
        lx = (len0 < len1)? len0 : len1;
    else
        lx = (lx < ly)? lx : ly;
    pct = 100.*(double)nm/(double)lx;
    fprintf(fx, "\n");
    fprintf(fx, "<%d match%s in an overlap of %d: %.2f percent similarity\n",
        nm, (nm == 1)? "" : "es", lx, pct);
    fprintf(fx, "<gaps in first sequence: %d", gapx);
    if (gapx) {
        (void) sprintf(outx, "(%d %s%s)",
            ngapx, (dna)? "base": "residue", (ngapx == 1)? "" : "s");
        fprintf(fx, "%s", outx);
    }
    fprintf(fx, ", gaps in second sequence: %d", gapy);
    if (gapy) {
        (void) sprintf(outx, "(%d %s%s)",
            ngapy, (dna)? "base": "residue", (ngapy == 1)? "" : "s");
        fprintf(fx, "%s", outx);
    }
    if (dna)
        fprintf(fx,
            "\n<score: %d (match = %d, mismatch = %d, gap penalty = %d + %d per base)\n",
            smax, DMAT, DMIS, DINS0, DINS1);
    else
        fprintf(fx,
            "\n<score: %d (Dayhoff PAM 250 matrix, gap penalty = %d + %d per residue)\n",
            smax, PINS0, PINS1);
    if (endgaps)
        fprintf(fx,
            "<endgaps penalized. left endgap: %d %s%s, right endgap: %d %s%s\n",
            firstgap, (dna)? "base": "residue", (firstgap == 1)? "" : "s",
            lastgap, (dna)? "base": "residue", (lastgap == 1)? "" : "s");
    else
        fprintf(fx, "<endgaps not penalized\n");
}
static      nm;                                                    /* matches in core -- for checking */

```

...getmat

TABLE 1-continued

```

static      lmax;          /* lengths of stripped file names */
static      ij[2];         /* jmp index for a path */
static      nc[2];         /* number at start of current line */
static      ni[2];         /* current elem number -- for gapping */
static      siz[2];
static char  *ps[2];        /* ptr to current element */
static char  *po[2];        /* ptr to next output char slot */
static char  ouf[2][P_LINE]; /* output line */
static char  star[P_LINE];  /* set by stars() */
/*
 * print alignment of described in struct path pp[]
 */
static
pr_align()
{
    int      nn;            /* char count */
    int      more;
    register i;
    for (i = 0, lmax = 0; i < 2; i++) {
        nn = stripname(name[i]);
        if (nn > lmax)
            lmax = nn;
        nc[i] = 1;
        ni[i] = 1;
        siz[i] = ij[i] = 0;
        ps[i] = seqx[i];
        po[i] = ouf[i];
    }
    for (nn = nm = 0, more = 1; more;) {
        for (i = more = 0; i < 2; i++) {
            /*
             * do we have more of this sequence?
             */
            if (!*ps[i])
                continue;
            more++;
            if (pp[i].spc) { /* leading space */
                *po[i]++ = ' ';
                pp[i].spc--;
            }
            else if (siz[i]) { /* in a gap */
                *po[i]++ = '-';
                siz[i]--;
            }
            else { /* we're putting a seq element
                    */
                *po[i] = *ps[i];
                if (islower(*ps[i]))
                    *ps[i] = toupper(*ps[i]);
                po[i]++;
                ps[i]++;
            }
            /*
             * are we at next gap for this seq?
             */
            if (ni[i] == pp[i].x[ij[i]]) {
                /*
                 * we need to merge all gaps
                 * at this location
                 */
                siz[i] == pp[i].n[ij[i]++];
                while (ni[i] == pp[i].x[ij[i]])
                    siz[i] += pp[i].n[ij[i]++];
            }
            ni[i]++;
        }
        if (++nn == olen || !more && nn) {
            dumpblock();
            for (i = 0; i < 2; i++)
                po[i] = ouf[i];
            nn = 0;
        }
    }
}
/*
 * dump a block of lines, including numbers, stars: pr_align()
 */
static
dumpblock()

```

pr_align

...pr_align

dumpblock

TABLE 1-continued

```

{
    register i;
    for(i = 0; i < 2; i++)
        *po[i]-- = '0';

    (void) putc('\n', fx);
    for (i = 0; i < 2; i++) {
        if (*out[i] && (*out[i] != ' ' || *(po[i]) != ' ')) {
            if (i == 0)
                nums(i);
            if (i == 0 && *out[1])
                stars();
            putline(i);
            if (i == 0 && *out[1])
                fprintf(fx, star);
            if (i == 1)
                nums(i);
        }
    }
}

/* put out a number line: dumpblock()
*/
static
nums(ix)
int    ix;    /* index in out[] holding seq line */
{
    char    nline[P__LINE];
    register i, j;
    register char *pn, *px, *py;
    for(pn = nline, i = 0; i < lmax+P__SPC; i++, pn++)
        *pn = ' ';
    for (i = nc[ix], py = out[ix]; *py; py++, pn++) {
        if (*py == ' ' || *py == '-');
        *pn = ' ';
        else {
            if (i%10 == 0 || (i == 1 && nc[ix] != 1)) {
                j = (i < 0)? -i : i;
                for (px = pn; j/= 10, px--)
                    *px = j%10 + '0';
                if (i < 0)
                    *px = '-';
            }
            else
                *pn = ' ';
            i++;
        }
    }
    *pn = '\0';
    nc[ix] = i;
    for (pn = nline; *pn; pn++)
        (void) putc(*pn, fx);
    (void) putc('\n', fx);
}

/*
 * put out a line (name, [num], seq. [num]): dumpblock()
 */
static
putline(ix)
int    ix;
{
    int    i;
    register char *px;
    for (px = name[ix], i = 0; *px && *px != ':'; px++, i++)
        (void) putc(*px, fx);
    for (; i < lmax+P__SPC; i++)
        (void) putc(' ', fx);
    /* these count from 1:
     * nl[] is current element (from 1)
     * nc[] is number at start of current line
     */
    for (px = out[ix]; *px; px++)
        (void) putc(*px&0x7F, fx);
    (void) putc('\n', fx);
}

/*
 * put a line of stars (seqs always in out[0], out[1]): dumpblock()
 */
static

```

...dumpblock

nums

putline

...putline

TABLE 1-continued

```

stars()
{
    int        i;
    register char *p0, *p1, cx, *px;
    if (!(*out[0] || (*out[0] == ' ' && (*p0[0] == ' ') ||
        !(*out[1] || (*out[1] == ' ' && (*p0[1] == ' ')))
        return;
    px = star;
    for (i = lmax+P_SPC; i; i--)
        *px++ = ' ';
    for (p0 = out[0], p1 = out[1]; *p0 && *p1; p0++, p1++) {
        if (isalpha(*p0) && isalpha(*p1)) {
            if (xbm[*p0-'A'] & xbm[*p1-'A']) {
                cx = '.';
                nm++;
            }
            else if (!dna && _day[*p0-'A'][*p1-'A'] > 0)
                cx = '.';
            else
                cx = ' ';
        }
        else
            cx = ' ';
        *px++ = cx;
    }
    *px++ = '\n';
    *px = '\0';
}
/*
 * strip path or prefix from pn, return len: pr_align()
 */
static
stripname(pn)
char *pn;          /* file name (may be path) */
{
    register char *px, *py;
    py = 0;
    for (px = pn; *px; px++)
        if (*px == '/')
            py = px + 1;
    if (py)
        (void) strcpy(pn, py);
    return(strlen(pn));
}
/*
 * cleanup() -- cleanup any tmp file
 * getseq() -- read in seq, set dna, len, maxlen
 * g_calloc() -- calloc() with error checkin
 * readjumps() -- get the good jumps, from tmp file if necessary
 * writejumps() -- write a filled array of jumps to a tmp file: nw()
 */
#include "nw.h"
#include <sys/file.h>
char *jname = "/tmp/homgXXXXXX"; /* tmp file for jumps */
FILE *fj;
int cleanup(); /* cleanup tmp file */
long lseek();
/*
 * remove any tmp file if we blow
 */
cleanup(i)
int i;
{
    if (fj)
        (void) unlink(jname);
    exit(i);
}
/*
 * read, return ptr to seq, set dna, len, maxlen
 * skip lines starting with ';', '<', or '>'
 * seq in upper or lower case
 */
char *
getseq(file, len)
char *file; /* file name */
int *len; /* seq len */
{
    char line[1024], *pseq;
    register char *px, *py;

```

stars

stripname

cleanup

getseq

TABLE 1-continued

```

int          natgc, tlen;
FILE         *fp;
if ((fp = fopen(file, "r")) == 0) {
    fprintf(stderr, "%s: can't read %s\n", prog, file);
    exit(1);
}
tlen = natgc = 0;
while (fgets(line, 1024, fp)) {
    if (*line == ';' || *line == '<' || *line == '>')
        continue;
    for (px = line; *px != '\n'; px++)
        if (isupper(*px) || islower(*px))
            tlen++;
}
if ((pseq = malloc((unsigned)(tlen+6))) == 0) {
    fprintf(stderr, "%s: malloc() failed to get %d bytes for %s\n", prog, tlen+6, file);
    exit(1);
}
pseq[0] = pseq[1] = pseq[2] = pseq[3] = '\0';

py = pseq + 4;
*len = tlen;
rewind(fp);
while (fgets(line, 1024, fp)) {
    if (*line == ';' || *line == '<' || *line == '>')
        continue;
    for (px = line; *px != '\n'; px++) {
        if (isupper(*px))
            *py++ = *px;
        else if (islower(*px))
            *py++ = toupper(*px);
        if (index("ATGCU", *(py-1)))
            natgc++;
    }
    *py++ = '\0';
    *py = '\0';
    (void) fclose(fp);
    dna = natgc > (tlen/3);
    return(pseq+4);
}

char *
g__calloc(msg, nx, sz)
char *msg;          /* program, calling routine */
int nx, sz;         /* number and size of elements */
{
    char *px, *calloc();
    if ((px = calloc((unsigned)nx, (unsigned)sz)) == 0) {
        if (*msg) {
            fprintf(stderr, "%s: g__calloc() failed %s (n= %d, sz= %d)\n", prog, msg, nx, sz);
            exit(1);
        }
    }
    return(px);
}

/*
 * get final jmps from dx[] or tmp file, set pp[], reset dmax: main()
 */
readjmps()
{
    int fd = -1;
    int siz, i0, i1;
    register i, j, xx;
    if (!f) {
        (void) fclose(f);
        if ((fd = open(jname, O_RDONLY, 0)) < 0) {
            fprintf(stderr, "%s: can't open() %s\n", prog, jname);
            cleanup(1);
        }
    }
    for (i = i0 = i1 = 0, dmax0 = dmax, xx = len0; i++;) {
        while (1) {
            for (j = dx[dmax].ijmp; j >= 0 && dx[dmax].jp.x[j] >= xx; j--)
                ;
            if (j < 0 && dx[dmax].offset && f) {
                (void) lseek(fd, dx[dmax].offset, 0);
                (void) read(fd, (char *)&dx[dmax].jp, sizeof(struct jmp));
                (void) read(fd, (char *)&dx[dmax].offset, sizeof(dx[dmax].offset));
            }
        }
    }
}

```

...getseq

g__calloc

readjmps

...readjmps

TABLE 1-continued

```

        dx[dmax].ijmp = MAXJMP-1;
    }
    else
        break;
    }
    if (i >= JMPS) {
        fprintf(stderr, "%s: too many gaps in alignment\n", prog);
        cleanup(1);
    }
    if (j >= 0) {
        siz = dx[dmax].jp.n[j];
        xx = dx[dmax].jp.x[j];
        dmax += siz;
        if (siz < 0) { /* gap in second seq */
            pp[1].n[i1] = -siz;
            xx += siz;
            /* id = xx - yy + len1 - 1
            */
            pp[1].x[i1] = xx - dmax + len1 - 1;
            gapy++;
            ngapy -= siz;
/* ignore MAXGAP when doing endgaps */
            siz = (-siz < MAXGAP || endgaps)? -siz : MAXGAP;
            i1++;
        }
        else if (siz > 0) { /* gap in first seq */
            pp[0].n[i0] = siz;
            pp[0].x[i0] = xx;
            gapx++;
            ngapx += siz;
/* ignore MAXGAP when doing endgaps */
            siz = (siz < MAXGAP || endgaps)? siz : MAXGAP;
            i0++;
        }
    }
    else
        break;
}
/* reverse the order of jmps
*/
for (j = 0, i0--, j < i0; j++, i0--) {
    i = pp[0].n[j]; pp[0].n[j] = pp[0].n[i0]; pp[0].n[i0] = i;
    i = pp[0].x[j]; pp[0].x[j] = pp[0].x[i0]; pp[0].x[i0] = i;
}
for (j = 0, i1--, j < i1; j++, i1--) {
    i = pp[1].n[j]; pp[1].n[j] = pp[1].n[i1]; pp[1].n[i1] = i;
    i = pp[1].x[j]; pp[1].x[j] = pp[1].x[i1]; pp[1].x[i1] = i;
}
if (fd >= 0)
    (void) close(fd);
if (fj) {
    (void) unlink(jname);
    fj = 0;
    offset = 0;
}
}
/*
 * write a filled jmp struct offset of the prev one (if any): nw()
 */
writejumps(ix)
    int    ix;
{
    char    *mktemp();
    if (!fj) {
        if (mktemp(jname) < 0) {
            fprintf(stderr, "%s: can't mktemp() %s\n", prog, jname);
            cleanup(1);
        }
        if ((fj = fopen(jname, "w")) == 0) {
            fprintf(stderr, "%s: can't write %s\n", prog, jname);
            exit(1);
        }
    }
    (void) fwrite((char *)&dx[ix].jp, sizeof(struct jmp), 1, fj);
    (void) fwrite((char *)&dx[ix].offset, sizeof(dx[ix].offset), 1, fj);
}

```

writejumps

TABLE 2

PRO	XXXXXXXXXXXXXXXX	(Length = 15 amino acids)
Comparison Protein	XXXXXXYYYYYYY	(Length = 12 amino acids)

% amino acid sequence identity=(the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide)=5 divided by 15=33.3%

TABLE 3

PRO	XXXXXXXXXX	(Length = 10 amino acids)
Comparison Protein	XXXXXXYYYYZZYZ	(Length = 15 amino acids)

% amino acid sequence identity=(the number of identically matching amino acid residues between the two polypeptide sequences as determined by ALIGN-2) divided by (the total number of amino acid residues of the PRO polypeptide)=5 divided by 10=50%

TABLE 4

PRO-DNA	NNNNNNNNNNNNNN	(Length = 14 nucleotides)
Comparison DNA	NNNNNNLLLLLLLL	(Length = 16 nucleotides)

% nucleic acid sequence identity=(the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence)=6 divided by 14=42.9%

TABLE 5

PRO-DNA	NNNNNNNNNNNN	(Length = 12 nucleotides)
Comparison DNA	NNNNLLLVV	(Length = 9 nucleotides)

% nucleic acid sequence identity=(the number of identically matching nucleotides between the two nucleic acid sequences as determined by ALIGN-2) divided by (the total number of nucleotides of the PRO-DNA nucleic acid sequence)=4 divided by 12=33.3%

II. Compositions and Methods of the Invention

A. Full-length PRO19598 Polypeptides

The present invention provides newly identified and isolated nucleotide sequences encoding polypeptides referred to in the present application as PRO19598 polypeptides. In particular, cDNAs encoding PRO19598 polypeptides have been identified and isolated, as disclosed in further detail in the Examples below. It is noted that proteins produced in separate expression rounds may be given different PRO numbers but the UNQ number is unique for any given DNA and the encoded protein, and will not be changed. However, for sake of simplicity, in the present specification the protein encoded by the full length native nucleic acid molecules disclosed herein as well as all further native homologues and variants included in the foregoing definition of PRO, will be referred to as "PRO/number", regardless of their origin or mode of preparation.

As disclosed in the Examples below, cDNA clones have been deposited with the ATCC. The actual nucleotide sequences of those clones can readily be determined by the skilled artisan by sequencing of the deposited clone using

routine methods in the art. The predicted amino acid sequence can be determined from the nucleotide sequence using routine skill. For the PRO19598 polypeptides and encoding nucleic acids described herein, Applicants have identified what is believed to be the reading frame best identifiable with the sequence information available at the time.

Using the ALIGN-2 sequence alignment computer program referenced above, it has been found that the full-length native sequence PRO19598 (shown in FIG. 2 and SEQ ID NO:2) has certain amino acid sequence identity with AF184971_1. Accordingly, it is presently believed that the PRO19598 polypeptide disclosed in the present application is a newly identified member of the cytokine receptor protein family and may possess one or more biological and/or immunological activities or properties typical of that protein family.

B. PRO19598 Polypeptide Variants

In addition to the full-length native sequence PRO19598 polypeptides described herein, it is contemplated that PRO19598 variants can be prepared. PRO19598 variants can be prepared by introducing appropriate nucleotide changes into the PRO19598 DNA, and/or by synthesis of the desired PRO19598 polypeptide. Those skilled in the art will appreciate that amino acid changes may alter post-translational processes of the PRO19598, such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

Variations in the native full-length sequence PRO19598 or in various domains of the PRO19598 described herein, can be made, for example, using any of the techniques and guidelines for conservative and non-conservative mutations set forth, for instance, in U.S. Pat. No. 5,364,934. Variations may be a substitution, deletion or insertion of one or more codons encoding the PRO19598 that results in a change in the amino acid sequence of the PRO19598 as compared with the native sequence PRO19598. Optionally the variation is by substitution of at least one amino acid with any other amino acid in one or more of the domains of the PRO19598. Guidance in determining which amino acid residue may be inserted, substituted or deleted without adversely affecting the desired activity may be found by comparing the sequence of the PRO19598 with that of homologous known protein molecules and minimizing the number of amino acid sequence changes made in regions of high homology. Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, i.e., conservative amino acid replacements. Insertions or deletions may optionally be in the range of about 1 to 5 amino acids. The variation allowed may be determined by systematically making insertions, deletions or substitutions of amino acids in the sequence and testing the resulting variants for activity exhibited by the full-length or mature native sequence.

PRO19598 polypeptide fragments are provided herein. Such fragments may be truncated at the N-terminus or C-terminus, or may lack internal residues, for example, when compared with a full length native protein. Certain fragments lack amino acid residues that are not essential for a desired biological activity of the PRO19598 polypeptide.

PRO19598 fragments may be prepared by any of a number of conventional techniques. Desired peptide fragments may be chemically synthesized. An alternative approach involves generating PRO19598 fragments by enzymatic digestion, e.g., by treating the protein with an enzyme known to cleave proteins at sites defined by par-

ticular amino acid residues, or by digesting the DNA with suitable restriction enzymes and isolating the desired fragment. Yet another suitable technique involves isolating and amplifying a DNA fragment encoding a desired polypeptide fragment, by polymerase chain reaction (PCR). Oligonucleotides that define the desired termini of the DNA fragment are employed at the 5' and 3' primers in the PCR. Preferably, PRO19598 polypeptide fragments share at least one biological and/or immunological activity with the native PRO19598 polypeptide disclosed herein.

In particular embodiments, conservative substitutions of interest are shown in Table 6 under the heading of preferred substitutions. If such substitutions result in a change in biological activity, then more substantial changes, denominated exemplary substitutions in Table 6, or as further described below in reference to amino acid classes, are introduced and the products screened.

TABLE 6

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	val; leu; ile	val
Arg (R)	lys; gln; asn	lys
Asn (N)	gln; his; lys; arg	gln
Asp (D)	glu	glu
Cys (C)	ser	ser
Gln (Q)	asn	asn
Glu (E)	asp	asp
Gly (G)	pro; ala	ala
His (H)	asn; gln; lys; arg	arg
Ile (I)	leu; val; met; ala; phe; norleucine	leu
Leu (L)	norleucine; ile; val; met; ala; phe	ile
Lys (K)	arg; gln; asn	arg
Met (M)	leu; phe; ile	leu
Phe (F)	leu; val; ile; ala; tyr	leu
Pro (P)	ala	ala
Ser (S)	thr	thr
Thr (T)	ser	ser
Trp (W)	tyr; phe	tyr
Tyr (Y)	trp; phe; thr; ser	phe
Val (V)	ile; leu; met; phe; ala; norleucine	leu

Substantial modifications in function or immunological identity of the PRO19598 polypeptide are accomplished by selecting substitutions that differ significantly in their effect on maintaining (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation, (b) the charge or hydrophobicity of the molecule at the target site, or (c) the bulk of the side chain. Naturally occurring residues are divided into groups based on common side-chain properties:

- (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- (2) neutral hydrophilic: cys, ser, thr;
- (3) acidic: asp, glu;
- (4) basic: asn, gln, his, lys, arg;
- (5) residues that influence chain orientation: gly, pro; and
- (6) aromatic: trp, tyr, phe.

Non-conservative substitutions will entail exchanging a member of one of these classes for another class. Such substituted residues also may be introduced into the conservative substitution sites or, more preferably, into the remaining (non-conserved) sites.

The variations can be made using methods known in the art such as oligonucleotide-mediated (site-directed) mutagenesis, alanine scanning, and PCR mutagenesis. Site-directed mutagenesis [Carter et al., *Nucl. Acids Res.*, 13:4331 (1986); Zoller et al., *Nucl. Acids Res.*, 10:6487

(1987)], cassette mutagenesis [Wells et al., *Gene*, 34:315 (1985)], restriction selection mutagenesis [Wells et al., *Philos. Trans. R. Soc. London SerA*, 317:415 (1986)] or other known techniques can be performed on the cloned DNA to produce the PRO19598 variant DNA.

Scanning amino acid analysis can also be employed to identify one or more amino acids along a contiguous sequence. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and cysteine. Alanine is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main-chain conformation of the variant [Cunningham and Wells, *Science*, 244: 1081-1085 (1989)]. Alanine is also typically preferred because it is the most common amino acid. Further, it is frequently found in both buried and exposed positions [Creighton, *The Proteins*, (W. H. Freeman & Co., N.Y.); Chothia, *J. Mol. Biol.*, 150:1 (1976)]. If alanine substitution does not yield adequate amounts of variant, an isoteric amino acid can be used.

C. Modifications of PRO19598

Covalent modifications of PRO19598 are included within the scope of this invention. One type of covalent modification includes reacting targeted amino acid residues of a PRO19598 polypeptide with an organic derivatizing agent that is capable of reacting with selected side chains or the N- or C-terminal residues of the PRO19598. Derivatization with bifunctional agents is useful, for instance, for crosslinking PRO19598 to a water-insoluble support matrix or surface for use in the method for purifying anti-PRO19598 antibodies, and vice-versa. Commonly used crosslinking agents include, e.g., 1,1-bis(diazoacetyl)-2-phenylethane, glutaraldehyde, N-hydroxysuccinimide esters, for example, esters with 4-azidosalicylic acid, homobifunctional imidoesters, including disuccinimidyl esters such as 3,3'-dithiobis(succinimidylpropionate), bifunctional maleimides such as bis-N-maleimido-1,8-octane and agents such as methyl-3-[(p-azidophenyl)dithio]propionimide.

Other modifications include amidation of glutamyl and asparaginy residues to the corresponding glutamyl and aspartyl residues, respectively, hydroxylation of proline and lysine, phosphorylation of hydroxyl groups of seryl or threonyl residues, methylation of the α -amino groups of lysine, arginine, and histidine side chains [T. E. Creighton, *Proteins: Structure and Molecular Properties*, W. H. Freeman & Co., San Francisco, pp. 79-86 (1983)], acetylation of the N-terminal amine, and amidation of any C-terminal carboxyl group.

Another type of covalent modification of the PRO19598 polypeptide included within the scope of this invention comprises altering the native glycosylation pattern of the polypeptide. "Altering the native glycosylation pattern" is intended for purposes herein to mean deleting one or more carbohydrate moieties found in native sequence PRO19598 (either by removing the underlying glycosylation site or by deleting the glycosylation by chemical and/or enzymatic means), and/or adding one or more glycosylation sites that are not present in the native sequence PRO19598. In addition, the phrase includes qualitative changes in the glycosylation of the native proteins, involving a change in the nature and proportions of the various carbohydrate moieties present.

Addition of glycosylation sites to the PRO19598 polypeptide may be accomplished by altering the amino acid sequence. The alteration may be made, for example, by the addition of, or substitution by, one or more serine or threonine residues to the native sequence PRO19598 (for

O-linked glycosylation sites). The PRO19598 amino acid sequence may optionally be altered through changes at the DNA level, particularly by mutating the DNA encoding the PRO19598 polypeptide at preselected bases such that codons are generated that will translate into the desired amino acids.

Another means of increasing the number of carbohydrate moieties on the PRO19598 polypeptide is by chemical or enzymatic coupling of glycosides to the polypeptide. Such methods are described in the art, e.g., in WO 87/05330 published Sep. 11, 1987, and in Aplin and Wriston, *CRC Crit. Rev. Biochem.*, pp. 259–306 (1981).

Removal of carbohydrate moieties present on the PRO19598 polypeptide may be accomplished chemically or enzymatically or by mutational substitution of codons encoding for amino acid residues that serve as targets for glycosylation. Chemical deglycosylation techniques are known in the art and described, for instance, by Hakimuddin, et al., *Arch. Biochem. Biophys.*, 259:52 (1987) and by Edge et al., *Anal. Biochem.*, 118:131 (1981). Enzymatic cleavage of carbohydrate moieties on polypeptides can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al., *Meth. Enzymol.*, 138:350 (1987).

Another type of covalent modification of PRO19598 comprises linking the PRO19598 polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Pat. Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

The PRO19598 of the present invention may also be modified in a way to form a chimeric molecule comprising PRO19598 fused to another, heterologous polypeptide or amino acid sequence.

In one embodiment, such a chimeric molecule comprises a fusion of the PRO19598 with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. The epitope tag is generally placed at the amino- or carboxyl-terminus of the PRO19598. The presence of such epitope-tagged forms of the PRO19598 can be detected using an antibody against the tag polypeptide. Also, provision of the epitope tag enables the PRO19598 to be readily purified by affinity purification using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag. Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-his) or poly-histidine-glycine (poly-his-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field et al., *Mol. Cell. Biol.*, 8:2159–2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan et al., *Molecular and Cellular Biology*, 5:3610–3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky et al., *Protein Engineering*, 3(6):547–553 (1990)]. Other tag polypeptides include the Flag-peptide [Hopp et al., *BioTechnology*, 6:1204–1210 (1988)]; the KT3 epitope peptide [Martin et al., *Science*, 255:192–194 (1992)]; an α -tubulin epitope peptide [Skinner et al., *J. Biol. Chem.*, 266:15163–15166 (1991)]; and the T7 gene 10 protein peptide tag [Lutz-Freyermuth et al., *Proc. Natl. Acad. Sci. USA*, 87:6393–6397 (1990)].

In an alternative embodiment, the chimeric molecule may comprise a fusion of the PRO19598 with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the chimeric molecule (also referred to as an “immunoadhesin”), such a fusion could be to the Fc region of an IgG molecule. The Ig fusions preferably include

the substitution of a soluble (transmembrane domain deleted or inactivated) form of a PRO19598 polypeptide in place of at least one variable region within an Ig molecule. In a particularly preferred embodiment, the immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG1 molecule. For the production of immunoglobulin fusions see also U.S. Pat. No. 5,428,130 issued Jun. 27, 1995.

D. Preparation of PRO19598

The description below relates primarily to production of PRO19598 by culturing cells transformed or transfected with a vector containing PRO19598 nucleic acid. It is, of course, contemplated that alternative methods, which are well known in the art, may be employed to prepare PRO19598. For instance, the PRO19598 sequence, or portions thereof, may be produced by direct peptide synthesis using solid-phase techniques [see, e.g., Stewart et al., *Solid-Phase Peptide Synthesis*, W. H. Freeman Co., San Francisco, Calif. (1969); Merrifield, *J. Am. Chem. Soc.*, 85:2149–2154 (1963)]. In vitro protein synthesis may be performed using manual techniques or by automation. Automated synthesis may be accomplished, for instance, using an Applied Biosystems Peptide Synthesizer (Foster City, Calif.) using manufacturer's instructions. Various portions of the PRO19598 may be chemically synthesized separately and combined using chemical or enzymatic methods to produce the full-length PRO19598.

1. Isolation of DNA Encoding PRO19598

DNA encoding PRO19598 may be obtained from a cDNA library prepared from tissue believed to possess the PRO19598 mRNA and to express it at a detectable level. Accordingly, human PRO19598 DNA can be conveniently obtained from a cDNA library prepared from human tissue, such as described in the Examples. The PRO19598-encoding gene may also be obtained from a genomic library or by known synthetic procedures (e.g., automated nucleic acid synthesis).

Libraries can be screened with probes (such as antibodies to the PRO19598 or oligonucleotides of at least about 20–80 bases) designed to identify the gene of interest or the protein encoded by it. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook et al., *Molecular Cloning: A Laboratory Manual* (New York: Cold Spring Harbor Laboratory Press, 1989). An alternative means to isolate the gene encoding PRO19598 is to use PCR methodology [Sambrook et al., supra; Dieffenbach et al., *PCR Primer: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1995)].

The Examples below describe techniques for screening a cDNA library. The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are minimized. The oligonucleotide is preferably labeled such that it can be detected upon hybridization to DNA in the library being screened. Methods of labeling are well known in the art, and include the use of radiolabels like ^{32}P -labeled ATP, biotinylation or enzyme labeling. Hybridization conditions, including moderate stringency and high stringency, are provided in Sambrook et al., supra.

Sequences identified in such library screening methods can be compared and aligned to other known sequences deposited and available in public databases such as GenBank or other private sequence databases. Sequence identity (at either the amino acid or nucleotide level) within defined regions of the molecule or across the full-length sequence can be determined using methods known in the art and as described herein.

Nucleic acid having protein coding sequence may be obtained by screening selected cDNA or genomic libraries using the deduced amino acid sequence disclosed herein for the first time, and, if necessary, using conventional primer extension procedures as described in Sambrook et al., supra, to detect precursors and processing intermediates of mRNA that may not have been reverse-transcribed into cDNA.

2. Selection and Transformation of Host Cells

Host cells are transfected or transformed with expression or cloning vectors described herein for PRO19598 production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH and the like, can be selected by the skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in *Mammalian Cell Biotechnology: a Practical Approach*, M. Butler, ed. (IRL Press, 1991) and Sambrook et al., supra.

Methods of eukaryotic cell transfection and prokaryotic cell transformation are known to the ordinarily skilled artisan, for example, CaCl_2 , CaPO_4 , liposome-mediated and electroporation. Depending on the host cell used, transformation is performed using standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in Sambrook et al., supra, or electroporation is generally used for prokaryotes. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw et al., *Gene*, 23:315 (1983) and WO 89/05859 published Jun. 29, 1989. For mammalian cells without such cell walls, the calcium phosphate precipitation method of Graham and van der Eb, *Virology*, 52:456-457 (1978) can be employed. General aspects of mammalian cell host system transfections have been described in U.S. Pat. No. 4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen et al., *J. Bact.*, 130:946 (1977) and Hsiao et al., *Proc. Natl. Acad. Sci. (USA)*, 76:3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, e.g., polybrene, polyornithine, may also be used. For various techniques for transforming mammalian cells, see Keown et al., *Methods in Enzymology*, 185:527-537 (1990) and Mansour et al., *Nature*, 336:348-352 (1988).

Suitable host cells for cloning or expressing the DNA in the vectors herein include prokaryote, yeast, or higher eukaryote cells. Suitable prokaryotes include but are not limited to eubacteria, such as Gram-negative or Gram-positive organisms, for example, Enterobacteriaceae such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 31,446); *E. coli* X1776 (ATCC 31,537); *E. coli* strain W3110 (ATCC 27,325) and K5 772 (ATCC 53,635). Other suitable prokaryotic host cells include Enterobacteriaceae such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as Bacilli such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41P disclosed in DD 266,710 published Apr. 12, 1989), *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. These examples are illustrative rather than limiting. Strain W3110 is one particularly preferred host or parent host because it is a common host strain for recombinant DNA product fermentations. Preferably, the host cell secretes minimal amounts of proteolytic enzymes. For example,

strain W3110 may be modified to effect a genetic mutation in the genes encoding proteins endogenous to the host, with examples of such hosts including *E. coli* W3110 strain 1A2, which has the complete genotype tonA; *E. coli* W3110 strain 9E4, which has the complete genotype tonA ptr3; *E. coli* W3110 strain 27C7 (ATCC 55,244), which has the complete genotype tonA ptr3 phoA E15 (argF-lac)169 degP ompT kan^r; *E. coli* W3110 strain 37D6, which has the complete genotype tonA ptr3 phoA E15 (argF-lac)169 degP ompT rbs7 ilvG kan^r; *E. coli* W3110 strain 40B4, which is strain 37D6 with a non-kanamycin resistant degP deletion mutation; and an *E. coli* strain having mutant periplasmic protease disclosed in U.S. Pat. No. 4,946,783 issued Aug. 7, 1990. Alternatively, in vitro methods of cloning, e.g., PCR or other nucleic acid polymerase reactions, are suitable.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts for PRO19598-encoding vectors. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism. Others include *Schizosaccharomyces pombe* (Beach and Nurse, *Nature*, 290:140 [1981]; EP 139,383 published May 2, 1985); *Kluyveromyces* hosts (U.S. Pat. No. 4,943,529; Fleer et al., *Bio/Technology*, 9:968-975 (1991)) such as, e.g., *K. lactis* (MW98-8C, CBS683, CBS4574; Louvencourt et al., *J. Bacteriol.*, 154(2):737-742 [1983]), *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickerhamii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilae* (ATCC 36,906; Van den Berg et al., *Bio/Technology*, 8:135 (1990)), *K. thermotolerans*, and *K. marxianus*; *yarrowia* (EP 402,226); *Pichia pastoris* (EP 183,070; Sreekrishna et al., *J. Basic Microbiol.*, 28:265-278 [1988]); *Candida*; *Trichodenna reesia* (EP 244,234); *Neurospora crassa* (Case et al., *Proc. Natl. Acad. Sci. USA*, 76:5259-5263 [1979]); *Schwanniomyces* such as *Schwanniomyces occidentalis* (EP 394,538 published Oct. 31, 1990); and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, *Tolypocladium* (WO 91/00357 published Jan. 10, 1991), and *Aspergillus* hosts such as *A. nidulans* (Ballance et al., *Biochem. Biophys. Res. Commun.*, 112:284-289 [1983]; Tilburn et al., *Gene*, 26:205-221 [1983]; Yelton et al., *Proc. Natl. Acad. Sci. USA*, 81:1470-1474 [1984]) and *A. niger* (Kelly and Hynes, *EMBO J.*, 4:475-479 [1985]). Methylotrophic yeasts are suitable herein and include, but are not limited to, yeast capable of growth on methanol selected from the genera consisting of *Hansenula*, *Candida*, *Kloeckera*, *Pichia*, *Saccharomyces*, *Torulopsis*, and *Rhodotorula*. A list of specific species that are exemplary of this class of yeasts may be found in C. Anthony, *The Biochemistry of Methylotrophs*, 269 (1982).

Suitable host cells for the expression of glycosylated PRO19598 are derived from multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila* S2 and *Spodoptera* Sf9, as well as plant cells. Examples of useful mammalian host cell lines include Chinese hamster ovary (CHO) and COS cells. More specific examples include monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for growth in suspension culture, Graham et al., *J. Gen. Virol.*, 36:59 (1977)); Chinese hamster ovary cells-DHFR (CHO, Urlaub and Chasin, *Proc. Natl. Acad. Sci. USA*, 77:4216 (1980)); mouse sertoli cells (TM4, Mather, *Biol. Reprod.*, 23:243-251 (1980)); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); and mouse mammary tumor (MMT 060562, ATCC CCL51). The selection of the appropriate host cell is deemed to be within the skill in the art.

3. Selection and Use of a Replicable Vector

The nucleic acid (e.g., cDNA or genomic DNA) encoding PRO19598 may be inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Various vectors are publicly available. The vector may, for example, be in the form of a plasmid, cosmid, viral particle, or phage. The appropriate nucleic acid sequence may be inserted into the vector by a variety of procedures. In general, DNA is inserted into an appropriate restriction endonuclease site(s) using techniques known in the art. Vector components generally include, but are not limited to, one or more of a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence. Construction of suitable vectors containing one or more of these components employs standard ligation techniques which are known to the skilled artisan.

The PRO19598 may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which may be a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. In general, the signal sequence may be a component of the vector, or it may be a part of the PRO19598-encoding DNA that is inserted into the vector. The signal sequence may be a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence may be, e.g., the yeast invertase leader, alpha factor leader (including *Saccharomyces* and *Kluyveromyces* α -factor leaders, the latter described in U.S. Pat. No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179 published Apr. 4, 1990), or the signal described in WO 90/13646 published Nov. 15, 1990. In mammalian cell expression, mammalian signal sequences may be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species, as well as viral secretory leaders.

Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2 μ plasmid origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells.

Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*.

An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the PRO19598-encoding nucleic acid, such as DHFR or thymidine kinase. An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared and propagated as described by Urlaub et al., *Proc. Natl. Acad. Sci. USA*, 77:4216 (1980). A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7 [Stinchcomb et al., *Nature*, 282:39 (1979); Kingsman et al., *Gene*, 7:141 (1979); Tschemper et al., *Gene*, 10:157 (1980)]. The *trp1* gene provides a selection marker for a mutant strain of yeast

lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 [Jones, *Genetics*, 85:12 (1977)].

Expression and cloning vectors usually contain a promoter operably linked to the PRO19598-encoding nucleic acid sequence to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well known. Promoters suitable for use with prokaryotic hosts include the β -lactamase and lactose promoter systems [Chang et al., *Nature*, 275:615 (1978); Goeddel et al., *Nature*, 281:544 (1979)], alkaline phosphatase, a tryptophan (*trp*) promoter system [Goeddel, *Nucleic Acids Res.*, 8:4057 (1980); EP 36,776], and hybrid promoters such as the *tac* promoter [deBoer et al., *Proc. Natl. Acad. Sci. USA*, 80:21-25 (1983)]. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding PRO19598.

Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase [Hitzeman et al., *J. Biol. Chem.*, 255:2073 (1980)] or other glycolytic enzymes [Hess et al., *J. Adv. Enzyme Reg.*, 7:149 (1968); Holland, *Biochemistry*, 17:4900 (1978)], such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657.

PRO19598 transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published Jul. 5, 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, and from heat-shock promoters, provided such promoters are compatible with the host cell systems.

Transcription of a DNA encoding the PRO19598 by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are cis-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin, α -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. The enhancer may be spliced into the vector at a position 5' or 3' to the PRO19598 coding sequence, but is preferably located at a site 5' from the promoter.

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of

eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated fragments in the untranslated portion of the mRNA encoding PRO19598.

Still other methods, vectors, and host cells suitable for adaptation to the synthesis of PRO19598 in recombinant vertebrate cell culture are described in Gething et al., *Nature*, 293:620-625 (1981); Mantei et al., *Nature*, 281:40-46 (1979); EP 117,060; and EP 117,058.

4. Detecting Gene Amplification/Expression

Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA [Thomas, *Proc. Natl. Acad. Sci. USA*, 77:5201-5205 (1980)], dot blotting (DNA analysis), or in situ hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

Gene expression, alternatively, may be measured by immunological methods, such as immunohistochemical staining of cells or tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any mammal. Conveniently, the antibodies may be prepared against a native sequence PRO19598 polypeptide or against a synthetic peptide based on the DNA sequences provided herein or against exogenous sequence fused to PRO19598 DNA and encoding a specific antibody epitope.

5. Purification of Polypeptide

Forms of PRO19598 may be recovered from culture medium or from host cell lysates. If membrane-bound, it can be released from the membrane using a suitable detergent solution (e.g. Triton-X 100) or by enzymatic cleavage. Cells employed in expression of PRO19598 can be disrupted by various physical or chemical means, such as freeze-thaw cycling, sonication, mechanical disruption, or cell lysing agents.

It may be desired to purify PRO19598 from recombinant cell proteins or polypeptides. The following procedures are exemplary of suitable purification procedures: by fractionation on an ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; protein A Sepharose columns to remove contaminants such as IgG; and metal chelating columns to bind epitope-tagged forms of the PRO19598. Various methods of protein purification may be employed and such methods are known in the art and described for example in Deutscher, *Methods in Enzymology*, 182 (1990); Scopes, *Protein Purification: Principles and Practice*, Springer-Verlag, New York (1982). The purification step(s) selected will depend, for example, on the nature of the production process used and the particular PRO19598 produced.

E. Uses for PRO19598

Nucleotide sequences (or their complement) encoding PRO19598 have various applications in the art of molecular biology, including uses as hybridization probes, in chromo-

some and gene mapping and in the generation of anti-sense RNA and DNA. PRO19598 nucleic acid will also be useful for the preparation of PRO19598 polypeptides by the recombinant techniques described herein.

The full-length native sequence PRO19598 gene, or portions thereof, may be used as hybridization probes for a cDNA library to isolate the full-length PRO19598 cDNA or to isolate still other cDNAs (for instance, those encoding naturally-occurring variants of PRO19598 or PRO19598 from other species) which have a desired sequence identity to the native PRO19598 sequence disclosed herein. Optionally, the length of the probes will be about 20 to about 50 bases. The hybridization probes may be derived from at least partially novel regions of the full length native nucleotide sequence wherein those regions may be determined without undue experimentation or from genomic sequences including promoters, enhancer elements and introns of native sequence PRO19598. By way of example, a screening method will comprise isolating the coding region of the PRO19598 gene using the known DNA sequence to synthesize a selected probe of about 40 bases. Hybridization probes may be labeled by a variety of labels, including radionucleotides such as ^{32}P or ^{35}S , or enzymatic labels such as alkaline phosphatase coupled to the probe via avidin/biotin coupling systems. Labeled probes having a sequence complementary to that of the PRO19598 gene of the present invention can be used to screen libraries of human cDNA, genomic DNA or mRNA to determine which members of such libraries the probe hybridizes to. Hybridization techniques are described in further detail in the Examples below.

Any EST sequences disclosed in the present application may similarly be employed as probes, using the methods disclosed herein.

Other useful fragments of the PRO19598 nucleic acids include antisense or sense oligonucleotides comprising a single-stranded nucleic acid sequence (either RNA or DNA) capable of binding to target PRO19598 mRNA (sense) or PRO19598 DNA (antisense) sequences. Antisense or sense oligonucleotides, according to the present invention, comprise a fragment of the coding region of PRO19598 DNA. Such a fragment generally comprises at least about 14 nucleotides, preferably from about 14 to 30 nucleotides. The ability to derive an antisense or a sense oligonucleotide, based upon a cDNA sequence encoding a given protein is described in, for example, Stein and Cohen (*Cancer Res.* 48:2659, 1988) and van der Krol et al. (*BioTechniques* 6:958, 1988).

Binding of antisense or sense oligonucleotides to target nucleic acid sequences results in the formation of duplexes that block transcription or translation of the target sequence by one of several means, including enhanced degradation of the duplexes, premature termination of transcription or translation, or by other means. The antisense oligonucleotides thus may be used to block expression of PRO19598 proteins. Antisense or sense oligonucleotides further comprise oligonucleotides having modified sugar-phosphodiester backbones (or other sugar linkages, such as those described in WO 91/06629) and wherein such sugar linkages are resistant to endogenous nucleases. Such oligonucleotides with resistant sugar linkages are stable in vivo (i.e., capable of resisting enzymatic degradation) but retain sequence specificity to be able to bind to target nucleotide sequences.

Other examples of sense or antisense oligonucleotides include those oligonucleotides which are covalently linked to organic moieties, such as those described in WO 90/10048, and other moieties that increases affinity of the

oligonucleotide for a target nucleic acid sequence, such as poly-(L-lysine). Further still, intercalating agents, such as ellipticine, and alkylating agents or metal complexes may be attached to sense or antisense oligonucleotides to modify binding specificities of the antisense or sense oligonucleotide for the target nucleotide sequence.

Antisense or sense oligonucleotides may be introduced into a cell containing the target nucleic acid sequence by any gene transfer method, including, for example, CaPO_4 -mediated DNA transfection, electroporation, or by using gene transfer vectors such as Epstein-Barr virus. In a preferred procedure, an antisense or sense oligonucleotide is inserted into a suitable retroviral vector. A cell containing the target nucleic acid sequence is contacted with the recombinant retroviral vector, either in vivo or ex vivo. Suitable retroviral vectors include, but are not limited to, those derived from the murine retrovirus M-MuLV, N2 (a retrovirus derived from M-MuLV), or the double copy vectors designated DCT5A, DCT5B and DCT5C (see WO 90/13641).

Sense or antisense oligonucleotides also may be introduced into a cell containing the target nucleotide sequence by formation of a conjugate with a ligand binding molecule, as described in WO 91/04753. Suitable ligand binding molecules include, but are not limited to, cell surface receptors, growth factors, other cytokines, or other ligands that bind to cell surface receptors. Preferably, conjugation of the ligand binding molecule does not substantially interfere with the ability of the ligand binding molecule to bind to its corresponding molecule or receptor, or block entry of the sense or antisense oligonucleotide or its conjugated version into the cell.

Alternatively, a sense or an antisense oligonucleotide may be introduced into a cell containing the target nucleic acid sequence by formation of an oligonucleotide-lipid complex, as described in WO 90/10448. The sense or antisense oligonucleotide-lipid complex is preferably dissociated within the cell by an endogenous lipase.

Antisense or sense RNA or DNA molecules are generally at least about 5 bases in length, about 10 bases in length, about 15 bases in length, about 20 bases in length, about 25 bases in length, about 30 bases in length, about 35 bases in length, about 40 bases in length, about 45 bases in length, about 50 bases in length, about 55 bases in length, about 60 bases in length, about 65 bases in length, about 70 bases in length, about 75 bases in length, about 80 bases in length, about 85 bases in length, about 90 bases in length, about 95 bases in length, about 100 bases in length, or more.

The probes may also be employed in PCR techniques to generate a pool of sequences for identification of closely related PRO19598 coding sequences.

Nucleotide sequences encoding a PRO19598 can also be used to construct hybridization probes for mapping the gene which encodes that PRO19598 and for the genetic analysis of individuals with genetic disorders. The nucleotide sequences provided herein may be mapped to a chromosome and specific regions of a chromosome using known techniques, such as in situ hybridization, linkage analysis against known chromosomal markers, and hybridization screening with libraries.

When the coding sequences for PRO19598 encode a protein which binds to another protein (example, where the PRO19598 is a receptor), the PRO19598 can be used in assays to identify the other proteins or molecules involved in the binding interaction. By such methods, inhibitors of the receptor/ligand binding interaction can be identified. Proteins involved in such binding interactions can also be used

to screen for peptide or small molecule inhibitors or agonists of the binding interaction. Also, the receptor PRO19598 can be used to isolate correlative ligand(s). Screening assays can be designed to find lead compounds that mimic the biological activity of a native PRO19598 or a receptor for PRO19598. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates. Small molecules contemplated include synthetic organic or inorganic compounds. The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays and cell based assays, which are well characterized in the art.

Nucleic acids which encode PRO19598 or its modified forms can also be used to generate either transgenic animals or "knock out" animals which, in turn, are useful in the development and screening of therapeutically useful reagents. A transgenic animal (e.g., a mouse or rat) is an animal having cells that contain a transgene, which transgene was introduced into the animal or an ancestor of the animal at a prenatal, e.g., an embryonic stage. A transgene is a DNA which is integrated into the genome of a cell from which a transgenic animal develops. In one embodiment, cDNA encoding PRO19598 can be used to clone genomic DNA encoding PRO19598 in accordance with established techniques and the genomic sequences used to generate transgenic animals that contain cells which express DNA encoding PRO19598. Methods for generating transgenic animals, particularly animals such as mice or rats, have become conventional in the art and are described, for example, in U.S. Pat. Nos. 4,736,866 and 4,870,009. Typically, particular cells would be targeted for PRO19598 transgene incorporation with tissue-specific enhancers. Transgenic animals that include a copy of a transgene encoding PRO19598 introduced into the germ line of the animal at an embryonic stage can be used to examine the effect of increased expression of DNA encoding PRO19598. Such animals can be used as tester animals for reagents thought to confer protection from, for example, pathological conditions associated with its overexpression. In accordance with this facet of the invention, an animal is treated with the reagent and a reduced incidence of the pathological condition, compared to untreated animals bearing the transgene, would indicate a potential therapeutic intervention for the pathological condition.

Alternatively, non-human homologues of PRO19598 can be used to construct a PRO19598 "knock out" animal which has a defective or altered gene encoding PRO19598 as a result of homologous recombination between the endogenous gene encoding PRO19598 and altered genomic DNA encoding PRO19598 introduced into an embryonic stem cell of the animal. For example, cDNA encoding PRO19598 can be used to clone genomic DNA encoding PRO19598 in accordance with established techniques. A portion of the genomic DNA encoding PRO19598 can be deleted or replaced with another gene, such as a gene encoding a selectable marker which can be used to monitor integration. Typically, several kilobases of unaltered flanking DNA (both at the 5' and 3' ends) are included in the vector [see e.g., Thomas and Capecchi, *Cell*, 51:503 (1987) for a description of homologous recombination vectors]. The vector is introduced into an embryonic stem cell line (e.g., by electroporation) and cells in which the introduced DNA has homologously recombined with the endogenous DNA are selected [see e.g., Li et al., *Cell*, 69:915 (1992)]. The selected cells are then injected into a blastocyst of an animal

(e.g., a mouse or rat) to form aggregation chimeras [see e.g., Bradley, in *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, E. J. Robertson, ed. (IRL, Oxford, 1987), pp. 113–152]. A chimeric embryo can then be implanted into a suitable pseudopregnant female foster animal and the embryo brought to term to create a “knock out” animal. Progeny harboring the homologously recombined DNA in their germ cells can be identified by standard techniques and used to breed animals in which all cells of the animal contain the homologously recombined DNA. Knock-out animals can be characterized for instance, for their ability to defend against certain pathological conditions and for their development of pathological conditions due to absence of the PRO19598 polypeptide.

Nucleic acid encoding the PRO19598 polypeptides may also be used in gene therapy. In gene therapy applications, genes are introduced into cells in order to achieve in vivo synthesis of a therapeutically effective genetic product, for example for replacement of a defective gene. “Gene therapy” includes both conventional gene therapy where a lasting effect is achieved by a single treatment, and the administration of gene therapeutic agents, which involves the one time or repeated administration of a therapeutically effective DNA or mRNA. Antisense RNAs and DNAs can be used as therapeutic agents for blocking the expression of certain genes in vivo. It has already been shown that short antisense oligonucleotides can be imported into cells where they act as inhibitors, despite their low intracellular concentrations caused by their restricted uptake by the cell membrane. (Zamecnik et al., *Proc. Natl. Acad. Sci. USA* 83:4143–4146 [1986]). The oligonucleotides can be modified to enhance their uptake, e.g. by substituting their negatively charged phosphodiester groups by uncharged groups.

There are a variety of techniques available for introducing nucleic acids into viable cells. The techniques vary depending upon whether the nucleic acid is transferred into cultured cells in vitro, or in vivo in the cells of the intended host. Techniques suitable for the transfer of nucleic acid into mammalian cells in vitro include the use of liposomes, electroporation, microinjection, cell fusion, DEAE-dextran, the calcium phosphate precipitation method, etc. The currently preferred in vivo gene transfer techniques include transfection with viral (typically retroviral) vectors and viral coat protein-liposome mediated transfection (Dzau et al., *Trends in Biotechnology* 11, 205–210 [1993]). In some situations it is desirable to provide the nucleic acid source with an agent that targets the target cells, such as an antibody specific for a cell surface membrane protein or the target cell, a ligand for a receptor on the target cell, etc. Where liposomes are employed, proteins which bind to a cell surface membrane protein associated with endocytosis may be used for targeting and/or to facilitate uptake, e.g. capsid proteins or fragments thereof tropic for a particular cell type, antibodies for proteins which undergo internalization in cycling, proteins that target intracellular localization and enhance intracellular half-life. The technique of receptor-mediated endocytosis is described, for example, by Wu et al., *J. Biol. Chem.* 262, 4429–4432 (1987); and Wagner et al., *Proc. Natl. Acad. Sci. USA* 87, 3410–3414 (1990). For review of gene marking and gene therapy protocols see Anderson et al., *Science* 256, 808–813 (1992).

The PRO19598 polypeptides described herein may also be employed as molecular weight markers for protein electrophoresis purposes and the isolated nucleic acid sequences may be used for recombinantly expressing those markers.

The nucleic acid molecules encoding the PRO19598 polypeptides or fragments thereof described herein are use-

ful for chromosome identification. In this regard, there exists an ongoing need to identify new chromosome markers, since relatively few chromosome marking reagents, based upon actual sequence data are presently available. Each PRO19598 nucleic acid molecule of the present invention can be used as a chromosome marker.

The PRO19598 polypeptides and nucleic acid molecules of the present invention may also be used diagnostically for tissue typing, wherein the PRO19598 polypeptides of the present invention may be differentially expressed in one tissue as compared to another, preferably in a diseased tissue as compared to a normal tissue of the same tissue type. PRO19598 nucleic acid molecules will find use for generating probes for PCR, Northern analysis, Southern analysis and Western analysis.

The PRO19598 polypeptides described herein may also be employed as therapeutic agents. The PRO19598 polypeptides of the present invention can be formulated according to known methods to prepare pharmaceutically useful compositions, whereby the PRO19598 product hereof is combined in admixture with a pharmaceutically acceptable carrier vehicle. Therapeutic formulations are prepared for storage by mixing the active ingredient having the desired degree of purity with optional physiologically acceptable carriers, excipients or stabilizers (*Remington's Pharmaceutical Sciences* 16th edition, Osol, A. Ed. (1980)), in the form of lyophilized formulations or aqueous solutions. Acceptable carriers, excipients or stabilizers are nontoxic to recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone, amino acids such as glycine, glutamine, asparagine, arginine or lysine; monosaccharides, disaccharides and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN™, PLURONICS™ or PEG.

The formulations to be used for in vivo administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes, prior to or following lyophilization and reconstitution.

Therapeutic compositions herein generally are placed into a container having a sterile access port, for example, an intravenous solution bag or vial having a stopper pierceable by a hypodermic injection needle.

The route of administration is in accord with known methods, e.g. injection or infusion by intravenous, intraperitoneal, intracerebral, intramuscular, intraocular, intraarterial or intralesional routes, topical administration, or by sustained release systems.

Dosages and desired drug concentrations of pharmaceutical compositions of the present invention may vary depending on the particular use envisioned. The determination of the appropriate dosage or route of administration is well within the skill of an ordinary physician. Animal experiments provide reliable guidance for the determination of effective doses for human therapy. Interspecies scaling of effective doses can be performed following the principles laid down by Mordenti, J. and Chappell, W. “The use of interspecies scaling in toxicokinetics” In *Toxicokinetics and New Drug Development*, Yacobi et al., Eds., Pergamon Press, New York 1989, pp. 42–96.

When in vivo administration of a PRO19598 polypeptide or agonist or antagonist thereof is employed, normal dosage

amounts may vary from about 10 ng/kg to up to 100 mg/kg of mammal body weight or more per day, preferably about 1 μ g/kg/day to 10 mg/kg/day, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature; see, for example, U.S. Pat. Nos. 4,657,760; 5,206,344; or 5,225,212. It is anticipated that different formulations will be effective for different treatment compounds and different disorders, that administration targeting one organ or tissue, for example, may necessitate delivery in a manner different from that to another organ or tissue.

Where sustained-release administration of a PRO19598 polypeptide is desired in a formulation with release characteristics suitable for the treatment of any disease or disorder requiring administration of the PRO19598 polypeptide, microencapsulation of the PRO19598 polypeptide is contemplated. Microencapsulation of recombinant proteins for sustained release has been successfully performed with human growth hormone (rhGH), interferon- (rhIFN-), interleukin-2, and MN rgp120. Johnson et al., *Nat. Med.*, 2:795-799 (1996); Yasuda, *Biomed. Ther.*, 27:1221-1223 (1993); Hora et al., *Bio/Technology*, 8:755-758 (1990); Cleland, "Design and Production of Single Immunization Vaccines Using Polylactide Polyglycolide Microsphere Systems," in *Vaccine Design: The Subunit and Adjuvant Approach*, Powell and Newman, eds, (Plenum Press: New York, 1995), pp. 439-462; WO 97/03692, WO 96/40072, WO 96/07399; and U.S. Pat. No. 5,654,010.

The sustained-release formulations of these proteins were developed using poly-lactic-co-glycolic acid (PLGA) polymer due to its biocompatibility and wide range of biodegradable properties. The degradation products of PLGA, lactic and glycolic acids, can be cleared quickly within the human body. Moreover, the degradability of this polymer can be adjusted from months to years depending on its molecular weight and composition. Lewis, "Controlled release of bioactive agents from lactide/glycolide polymer," in: M. Chasin and R. Langer (Eds.), *Biodegradable Polymers as Drug Delivery Systems* (Marcel Dekker: New York, 1990), pp. 1-41.

This invention encompasses methods of screening compounds to identify those that mimic the PRO19598 polypeptide (agonists) or prevent the effect of the PRO19598 polypeptide (antagonists). Screening assays for antagonist drug candidates are designed to identify compounds that bind or complex with the PRO19598 polypeptides encoded by the genes identified herein, or otherwise interfere with the interaction of the encoded polypeptides with other cellular proteins. Such screening assays will include assays amenable to high-throughput screening of chemical libraries, making them particularly suitable for identifying small molecule drug candidates.

The assays can be performed in a variety of formats, including protein-protein binding assays, biochemical screening assays, immunoassays, and cell-based assays, which are well characterized in the art.

All assays for antagonists are common in that they call for contacting the drug candidate with a PRO19598 polypeptide encoded by a nucleic acid identified herein under conditions and for a time sufficient to allow these two components to interact.

In binding assays, the interaction is binding and the complex formed can be isolated or detected in the reaction mixture. In a particular embodiment, the PRO19598 polypeptide encoded by the gene identified herein or the drug candidate is immobilized on a solid phase, e.g., on a microtiter plate, by covalent or non-covalent attachments.

Non-covalent attachment generally is accomplished by coating the solid surface with a solution of the PRO19598 polypeptide and drying. Alternatively, an immobilized antibody, e.g., a monoclonal antibody, specific for the PRO19598 polypeptide to be immobilized can be used to anchor it to a solid surface. The assay is performed by adding the non-immobilized component, which may be labeled by a detectable label, to the immobilized component, e.g., the coated surface containing the anchored component. When the reaction is complete, the non-reacted components are removed, e.g., by washing, and complexes anchored on the solid surface are detected. When the originally non-immobilized component carries a detectable label, the detection of label immobilized on the surface indicates that complexing occurred. Where the originally non-immobilized component does not carry a label, complexing can be detected, for example, by using a labeled antibody specifically binding the immobilized complex.

If the candidate compound interacts with but does not bind to a particular PRO19598 polypeptide encoded by a gene identified herein, its interaction with that polypeptide can be assayed by methods well known for detecting protein-protein interactions. Such assays include traditional approaches, such as, e.g., cross-linking, co-immunoprecipitation, and co-purification through gradients or chromatographic columns. In addition, protein-protein interactions can be monitored by using a yeast-based genetic system described by Fields and co-workers (Fields and Song, *Nature (London)*, 340:245-246 (1989); Chien et al., *Proc. Natl. Acad. Sci. USA*, 88:9578-9582 (1991)) as disclosed by Chevray and Nathans, *Proc. Natl. Acad. Sci. USA*, 89: 5789-5793 (1991). Many transcriptional activators, such as yeast GAL4, consist of two physically discrete modular domains, one acting as the DNA-binding domain, the other one functioning as the transcription-activation domain. The yeast expression system described in the foregoing publications (generally referred to as the "two-hybrid system") takes advantage of this property, and employs two hybrid proteins, one in which the target protein is fused to the DNA-binding domain of GAL4, and another, in which candidate activating proteins are fused to the activation domain. The expression of a GAL1-lacZ reporter gene under control of a GAL4-activated promoter depends on reconstitution of GAL4 activity via protein-protein interaction. Colonies containing interacting polypeptides are detected with a chromogenic substrate for β -galactosidase. A complete kit (MATCHMAKERTM) for identifying protein-protein interactions between two specific proteins using the two-hybrid technique is commercially available from Clontech. This system can also be extended to map protein domains involved in specific protein interactions as well as to pinpoint amino acid residues that are crucial for these interactions.

Compounds that interfere with the interaction of a gene encoding a PRO19598 polypeptide identified herein and other intra- or extracellular components can be tested as follows: usually a reaction mixture is prepared containing the product of the gene and the intra- or extracellular component under conditions and for a time allowing for the interaction and binding of the two products. To test the ability of a candidate compound to inhibit binding, the reaction is run in the absence and in the presence of the test compound. In addition, a placebo may be added to a third reaction mixture, to serve as positive control. The binding (complex formation) between the test compound and the intra- or extracellular component present in the mixture is monitored as described hereinabove. The formation of a

complex in the control reaction(s) but not in the reaction mixture containing the test compound indicates that the test compound interferes with the interaction of the test compound and its reaction partner.

To assay for antagonists, the PRO19598 polypeptide may be added to a cell along with the compound to be screened for a particular activity and the ability of the compound to inhibit the activity of interest in the presence of the PRO19598 polypeptide indicates that the compound is an antagonist to the PRO19598 polypeptide. Alternatively, antagonists may be detected by combining the PRO19598 polypeptide and a potential antagonist with membrane-bound PRO19598 polypeptide receptors or recombinant receptors under appropriate conditions for a competitive inhibition assay. The PRO19598 polypeptide can be labeled, such as by radioactivity, such that the number of PRO19598 polypeptide molecules bound to the receptor can be used to determine the effectiveness of the potential antagonist. The gene encoding the receptor can be identified by numerous methods known to those of skill in the art, for example, ligand panning and FACS sorting. Coligan et al., *Current Protocols in Immun.*, 1(2): Chapter 5 (1991). Preferably, expression cloning is employed wherein polyadenylated RNA is prepared from a cell responsive to the PRO19598 polypeptide and a cDNA library created from this RNA is divided into pools and used to transfect COS cells or other cells that are not responsive to the PRO19598 polypeptide. Transfected cells that are grown on glass slides are exposed to labeled PRO19598 polypeptide. The PRO19598 polypeptide can be labeled by a variety of means including iodination or inclusion of a recognition site for a site-specific protein kinase. Following fixation and incubation, the slides are subjected to autoradiographic analysis. Positive pools are identified and sub-pools are prepared and re-transfected using an interactive sub-pooling and re-screening process, eventually yielding a single clone that encodes the putative receptor.

As an alternative approach for receptor identification, labeled PRO19598 polypeptide can be photoaffinity-linked with cell membrane or extract preparations that express the receptor molecule. Cross-linked material is resolved by PAGE and exposed to X-ray film. The labeled complex containing the receptor can be excised, resolved into peptide fragments, and subjected to protein micro-sequencing. The amino acid sequence obtained from micro-sequencing would be used to design a set of degenerate oligonucleotide probes to screen a cDNA library to identify the gene encoding the putative receptor.

In another assay for antagonists, mammalian cells or a membrane preparation expressing the receptor would be incubated with labeled PRO19598 polypeptide in the presence of the candidate compound. The ability of the compound to enhance or block this interaction could then be measured.

More specific examples of potential antagonists include an oligonucleotide that binds to the fusions of immunoglobulin with PRO19598 polypeptide, and, in particular, antibodies including, without limitation, poly- and monoclonal antibodies and antibody fragments, single-chain antibodies, anti-idiotypic antibodies, and chimeric or humanized versions of such antibodies or fragments, as well as human antibodies and antibody fragments. Alternatively, a potential antagonist may be a closely related protein, for example, a mutated form of the PRO19598 polypeptide that recognizes the receptor but imparts no effect, thereby competitively inhibiting the action of the PRO19598 polypeptide.

Another potential PRO19598 polypeptide antagonist is an antisense RNA or DNA construct prepared using antisense

technology, where, e.g., an antisense RNA or DNA molecule acts to block directly the translation of mRNA by hybridizing to targeted mRNA and preventing protein translation. Antisense technology can be used to control gene expression through triple-helix formation or antisense DNA or RNA, both of which methods are based on binding of a polynucleotide to DNA or RNA. For example, the 5' coding portion of the polynucleotide sequence, which encodes the mature PRO19598 polypeptides herein, is used to design an antisense RNA oligonucleotide of from about 10 to 40 base pairs in length. A DNA oligonucleotide is designed to be complementary to a region of the gene involved in transcription (triple helix—see Lee et al., *Nucl. Acids Res.*, 6:3073 (1979); Cooney et al., *Science*, 241: 456 (1988); Dervan et al., *Science*, 251:1360 (1991)), thereby preventing transcription and the production of the PRO19598 polypeptide. The antisense RNA oligonucleotide hybridizes to the mRNA in vivo and blocks translation of the mRNA molecule into the PRO19598 polypeptide (antisense—Okano, *Neurochem.*, 56:560 (1991); *Oligodeoxynucleotides as Antisense Inhibitors of Gene Expression* (CRC Press: Boca Raton, Fla., 1988). The oligonucleotides described above can also be delivered to cells such that the antisense RNA or DNA may be expressed in vivo to inhibit production of the PRO19598 polypeptide. When antisense DNA is used, oligodeoxyribonucleotides derived from the translation-initiation site, e.g., between about -10 and +10 positions of the target gene nucleotide sequence, are preferred.

Potential antagonists include small molecules that bind to the active site, the receptor binding site, or growth factor or other relevant binding site of the PRO19598 polypeptide, thereby blocking the normal biological activity of the PRO19598 polypeptide. Examples of small molecules include, but are not limited to, small peptides or peptide-like molecules, preferably soluble peptides, and synthetic non-peptidyl organic or inorganic compounds.

Ribozymes are enzymatic RNA molecules capable of catalyzing the specific cleavage of RNA. Ribozymes act by sequence-specific hybridization to the complementary target RNA, followed by endonucleolytic cleavage. Specific ribozyme cleavage sites within a potential RNA target can be identified by known techniques. For further details see, e.g., Rossi, *Current Biology*, 4:469-471 (1994), and PCT publication No. WO 97/33551 (published Sep. 18, 1997).

Nucleic acid molecules in triple-helix formation used to inhibit transcription should be single-stranded and composed of deoxynucleotides. The base composition of these oligonucleotides is designed such that it promotes triple-helix formation via Hoogsteen base-pairing rules, which generally require sizeable stretches of purines or pyrimidines on one strand of a duplex. For further details see, e.g., PCT publication No. WO 97/33551, supra.

These small molecules can be identified by any one or more of the screening assays discussed hereinabove and/or by any other screening techniques well known for those skilled in the art.

Diagnostic and therapeutic uses of the herein disclosed molecules may also be based upon the positive functional assay hits disclosed and described below.

F. Anti-PRO19598 Antibodies

The present invention further provides anti-PRO19598 antibodies. Exemplary antibodies include polyclonal, monoclonal, humanized, bispecific, and heteroconjugate antibodies.

1. Polyclonal Antibodies

The anti-PRO19598 antibodies may comprise polyclonal antibodies. Methods of preparing polyclonal antibodies are

known to the skilled artisan. Polyclonal antibodies can be raised in a mammal, for example, by one or more injections of an immunizing agent and, if desired, an adjuvant. Typically, the immunizing agent and/or adjuvant will be injected in the mammal by multiple subcutaneous or intraperitoneal injections. The immunizing agent may include the PRO19598 polypeptide or a fusion protein thereof. It may be useful to conjugate the immunizing agent to a protein known to be immunogenic in the mammal being immunized. Examples of such immunogenic proteins include but are not limited to keyhole limpet hemocyanin, serum albumin, bovine thyroglobulin, and soybean trypsin inhibitor. Examples of adjuvants which may be employed include Freund's complete adjuvant and MPL-TDM adjuvant (monophosphoryl Lipid A, synthetic trehalose dicorynomycolate). The immunization protocol may be selected by one skilled in the art without undue experimentation.

2. Monoclonal Antibodies

The anti-PRO19598 antibodies may, alternatively, be monoclonal antibodies. Monoclonal antibodies may be prepared using hybridoma methods, such as those described by Kohler and Milstein, *Nature*, 256:495 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized in vitro.

The immunizing agent will typically include the PRO19598 polypeptide or a fusion protein thereof. Generally, either peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell [Goding, *Monoclonal Antibodies: Principles and Practice*, Academic Press, (1986) pp. 59-103]. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells may be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

Preferred immortalized cell lines are those that fuse efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, Calif. and the American Type Culture Collection, Manassas, Va. Human myeloma and mouse-human heteromyeloma cell lines also have been described for the production of human monoclonal antibodies [Kozbor, *J. Immunol.*, 133:3001 (1984); Brodeur et al., *Monoclonal Antibody Production Techniques and Applications*, Marcel Dekker, Inc., New York, (1987) pp. 51-63].

The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal

antibodies directed against PRO19598. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an in vitro binding assay, such as radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, *Anal. Biochem.*, 107:220 (1980).

After the desired hybridoma cells are identified, the clones may be subcloned by limiting dilution procedures and grown by standard methods [Goding, *supra*]. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium and RPMI-1640 medium. Alternatively, the hybridoma cells may be grown in vivo as ascites in a mammal.

The monoclonal antibodies secreted by the subclones may be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

The monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U.S. Pat. No. 4,816,567. DNA encoding the monoclonal antibodies of the invention can be readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes that are capable of binding specifically to genes encoding the heavy and light chains of murine antibodies). The hybridoma cells of the invention serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous murine sequences [U.S. Pat. No. 4,816,567; Morrison et al., *supra*] or by covalently joining to the immunoglobulin coding sequence all or part of the coding sequence for a non-immunoglobulin polypeptide. Such a non-immunoglobulin polypeptide can be substituted for the constant domains of an antibody of the invention, or can be substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody.

The antibodies may be monovalent antibodies. Methods for preparing monovalent antibodies are well known in the art. For example, one method involves recombinant expression of immunoglobulin light chain and modified heavy chain. The heavy chain is truncated generally at any point in the Fc region so as to prevent heavy chain crosslinking. Alternatively, the relevant cysteine residues are substituted with another amino acid residue or are deleted so as to prevent crosslinking.

In vitro methods are also suitable for preparing monovalent antibodies. Digestion of antibodies to produce fragments thereof, particularly, Fab fragments, can be accomplished using routine techniques known in the art.

3. Human and Humanized Antibodies

The anti-PRO19598 antibodies of the invention may further comprise humanized antibodies or human antibodies. Humanized forms of non-human (e.g., murine) antibodies are chimeric immunoglobulins, immunoglobulin chains or fragments thereof (such as Fv, Fab, Fab', F(ab')₂ or other antigen-binding subsequences of antibodies) which contain

minimal sequence derived from non-human immunoglobulin. Humanized antibodies include human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species (donor antibody) such as mouse, rat or rabbit having the desired specificity, affinity and capacity. In some instances, Fv framework residues of the human immunoglobulin are replaced by corresponding non-human residues. Humanized antibodies may also comprise residues which are found neither in the recipient antibody nor in the imported CDR or framework sequences. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region (Fc), typically that of a human immunoglobulin [Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-329 (1988); and Presta, *Curr. Op. Struct. Bio.*, 2:593-596 (1992)].

Methods for humanizing non-human antibodies are well known in the art. Generally, a humanized antibody has one or more amino acid residues introduced into it from a source which is non-human. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and co-workers [Jones et al., *Nature*, 321:522-525 (1986); Riechmann et al., *Nature*, 332:323-327 (1988); Verhoeven et al., *Science*, 239: 1534-1536 (1988)], by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody. Accordingly, such "humanized" antibodies are chimeric antibodies (U.S. Pat. No. 4,816,567), wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent antibodies.

Human antibodies can also be produced using various techniques known in the art, including phage display libraries [Hoogenboom and Winter, *J. Mol. Biol.*, 227:381 (1991); Marks et al., *J. Mol. Biol.*, 222:581 (1991)]. The techniques of Cole et al. and Boerner et al. are also available for the preparation of human monoclonal antibodies (Cole et al., *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, p. 77 (1985) and Boerner et al., *J. Immunol.*, 147(1):86-95 (1991)]. Similarly, human antibodies can be made by introducing of human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Pat. Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in the following scientific publications: Marks et al., *BioTechnology* 10, 779-783 (1992); Lonberg et al., *Nature* 368 856-859 (1994); Morrison, *Nature* 368, 812-13 (1994); Fishwild et al., *Nature Biotechnology* 14, 845-51 (1996); Neuberger, *Nature Biotechnology* 14, 826 (1996); Lonberg and Huszar, *Intern. Rev. Immunol.* 13 65-93 (1995).

The antibodies may also be affinity matured using known selection and/or mutagenesis methods as described above. Preferred affinity matured antibodies have an affinity which is five times, more preferably 10 times, even more preferably 20 or 30 times greater than the starting antibody (generally murine, humanized or human) from which the matured antibody is prepared.

4. Bispecific Antibodies

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least two different antigens. In the present case, one of the binding specificities is for the PRO19598, the other one is for any other antigen, and preferably for a cell-surface protein or receptor or receptor subunit.

Methods for making bispecific antibodies are known in the art. Traditionally, the recombinant production of bispecific antibodies is based on the co-expression of two immunoglobulin heavy-chain/light-chain pairs, where the two heavy chains have different specificities [Milstein and Cuello, *Nature*, 305:537-539 (1983)]. Because of the random assortment of immunoglobulin heavy and light chains, these hybridomas (quadromas) produce a potential mixture of ten different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule is usually accomplished by affinity chromatography steps. Similar procedures are disclosed in WO 93/08829, published May 13, 1993, and in Trautnecker et al., *EMBO J.*, 10:3655-3659 (1991).

Antibody variable domains with the desired binding specificities (antibody-antigen combining sites) can be fused to immunoglobulin constant domain sequences. The fusion preferably is with an immunoglobulin heavy-chain constant domain, comprising at least part of the hinge, CH2, and CH3 regions. It is preferred to have the first heavy-chain constant region (CH1) containing the site necessary for light-chain binding present in at least one of the fusions. DNAs encoding the immunoglobulin heavy-chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are co-transfected into a suitable host organism. For further details of generating bispecific antibodies see, for example, Suresh et al., *Methods in Enzymology*, 121:210 (1986).

According to another approach described in WO 96/27011, the interface between a pair of antibody molecules can be engineered to maximize the percentage of heterodimers which are recovered from recombinant cell culture. The preferred interface comprises at least a part of the CH3 region of an antibody constant domain. In this method, one or more small amino acid side chains from the interface of the first antibody molecule are replaced with larger side chains (e.g. tyrosine or tryptophan). Compensatory "cavities" of identical or similar size to the large side chain(s) are created on the interface of the second antibody molecule by replacing large amino acid side chains with smaller ones (e.g. alanine or threonine). This provides a mechanism for increasing the yield of the heterodimer over other unwanted end-products such as homodimers.

Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g. F(ab')₂ bispecific antibodies). Techniques for generating bispecific antibodies from antibody fragments have been described in the literature. For example, bispecific antibodies can be prepared can be prepared using chemical linkage. Brennan et al., *Science* 229:81 (1985) describe a procedure wherein intact antibodies are proteolytically cleaved to generate F(ab')₂ fragments. These fragments are reduced in the presence of the dithiol complexing agent sodium arsenite to stabilize vicinal dithi-

ols and prevent intermolecular disulfide formation. The Fab' fragments generated are then converted to thionitrobenzoate (TNB) derivatives. One of the Fab'-TNB derivatives is then reconverted to the Fab'-thiol by reduction with mercaptoethylamine and is mixed with an equimolar amount of the other Fab'-TNB derivative to form the bispecific antibody. The bispecific antibodies produced can be used as agents for the selective immobilization of enzymes.

Fab' fragments may be directly recovered from *E. coli* and chemically coupled to form bispecific antibodies. Shalaby et al., *J. Exp. Med.* 175:217-225 (1992) describe the production of a fully humanized bispecific antibody F(ab')₂ molecule. Each Fab' fragment was separately secreted from *E. coli* and subjected to directed chemical coupling in vitro to form the bispecific antibody. The bispecific antibody thus formed was able to bind to cells overexpressing the ErbB2 receptor and normal human T cells, as well as trigger the lytic activity of human cytotoxic lymphocytes against human breast tumor targets.

Various technique for making and isolating bispecific antibody fragments directly from recombinant cell culture have also been described. For example, bispecific antibodies have been produced using leucine zippers. Kostelny et al., *J. Immunol.* 148(5): 1547-1553 (1992). The leucine zipper peptides from the Fos and Jun proteins were linked to the Fab' portions of two different antibodies by gene fusion. The antibody homodimers were reduced at the hinge region to form monomers and then re-oxidized to form the antibody heterodimers. This method can also be utilized for the production of antibody homodimers. The "diabody" technology described by Hollinger et al., *Proc. Natl. Acad. Sci. USA* 90:6444-6448 (1993) has provided an alternative mechanism for making bispecific antibody fragments. The fragments comprise a heavy-chain variable domain (V_H) connected to a light-chain variable domain (V_L) by a linker which is too short to allow pairing between the two domains on the same chain. Accordingly, the V_H and V_L domains of one fragment are forced to pair with the complementary V_L and V_H domains of another fragment, thereby forming two antigen-binding sites. Another strategy for making bispecific antibody fragments by the use of single-chain Fv (sFv) dimers has also been reported. See, Gruber et al., *J. Immunol.* 152:5368 (1994). Antibodies with more than two valencies are contemplated. For example, trispecific antibodies can be prepared. Tutt et al., *J. Immunol.* 147:60 (1991).

Exemplary bispecific antibodies may bind to two different epitopes on a given PRO19598 polypeptide herein. Alternatively, an anti-PRO19598 polypeptide arm may be combined with an arm which binds to a triggering molecule on a leukocyte such as a T-cell receptor molecule (e.g. CD2, CD3, CD28, or B7), or Fc receptors for IgG (FcγR), such as FcγRI (CD64), FcγRII (CD32) and FcγRIII (CD16) so as to focus cellular defense mechanisms to the cell expressing the particular PRO19598 polypeptide. Bispecific antibodies may also be used to localize cytotoxic agents to cells which express a particular PRO19598 polypeptide. These antibodies possess a PRO19598-binding arm and an arm which binds a cytotoxic agent or a radionuclide chelator, such as EOTUBE, DPTA, DOTA, or TETA. Another bispecific antibody of interest binds the PRO19598 polypeptide and further binds tissue factor (TF).

5. Heteroconjugate Antibodies

Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells [U.S. Pat. No. 4,676,980], and for

treatment of HIV infection [WO 91/00360; WO 92/200373; EP 03089]. It is contemplated that the antibodies may be prepared in vitro using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose include iminothiolate and methyl-4-mercaptobutyrimidate and those disclosed, for example, in U.S. Pat. No. 4,676,980.

6. Effector Function Engineering

It may be desirable to modify the antibody of the invention with respect to effector function, so as to enhance, e.g., the effectiveness of the antibody in treating cancer. For example, cysteine residue(s) may be introduced into the Fc region, thereby allowing interchain disulfide bond formation in this region. The homodimeric antibody thus generated may have improved internalization capability and/or increased complement-mediated cell killing and antibody-dependent cellular cytotoxicity (ADCC). See Caron et al., *J. Exp. Med.*, 176: 1191-1195 (1992) and Shopes, *J. Immunol.*, 148: 2918-2922 (1992). Homodimeric antibodies with enhanced anti-tumor activity may also be prepared using heterobifunctional cross-linkers as described in Wolff et al. *Cancer Research*, 53: 2560-2565 (1993). Alternatively, an antibody can be engineered that has dual Fc regions and may thereby have enhanced complement lysis and ADCC capabilities. See Stevenson et al., *Anti-Cancer Drug Design*, 3: 219-230 (1989).

7. Immunoconjugates

The invention also pertains to immunoconjugates comprising an antibody conjugated to a cytotoxic agent such as a chemotherapeutic agent, toxin (e.g., an enzymatically active toxin of bacterial, fungal, plant, or animal origin, or fragments thereof), or a radioactive isotope (i.e., a radioconjugate).

Chemotherapeutic agents useful in the generation of such immunoconjugates have been described above. Enzymatically active toxins and fragments thereof that can be used include diphtheria A chain, nonbinding active fragments of diphtheria toxin, exotoxin A chain (from *Pseudomonas aeruginosa*), ricin A chain, abrin A chain, modeccin A chain, alpha-sarcin, *Aleurites fordii* proteins, dianthin proteins, *Phytolacca americana* proteins (PAPI, PAPII, and PAP-S), momordica charantia inhibitor, curcin, crotin, sapaonaria officinalis inhibitor, gelonin, mitogellin, restrictocin, phenomycin, enomycin, and the tricothecenes. A variety of radionuclides are available for the production of radioconjugated antibodies. Examples include ²¹²Bi, ¹³¹I, ¹³¹In, ⁹⁰Y, and ¹⁸⁶Re. Conjugates of the antibody and cytotoxic agent are made using a variety of bifunctional protein-coupling agents such as N-succinimidyl-3-(2-pyridyldithiol) propionate (SPDP), ininothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl)hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as tolyene 2,6-diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, a ricin immunotoxin can be prepared as described in Vitetta et al., *Science*, 238: 1098 (1987). Carbon-14-labeled 1-isothiocyanatobenzyl-3-methyldiethylene triaminepentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionuclide to the antibody. See WO94/11026.

In another embodiment, the antibody may be conjugated to a "receptor" (such streptavidin) for utilization in tumor

pretargeting wherein the antibody-receptor conjugate is administered to the patient, followed by removal of unbound conjugate from the circulation using a clearing agent and then administration of a "ligand" (e.g., avidin) that is conjugated to a cytotoxic agent (e.g., a radionucleotide).

8. Immunoliposomes

The antibodies disclosed herein may also be formulated as immunoliposomes. Liposomes containing the antibody are prepared by methods known in the art, such as described in Epstein et al., *Proc. Natl. Acad. Sci. USA*, 82: 3688 (1985); Hwang et al., *Proc. Natl. Acad. Sci. USA*, 77: 4030 (1980); and U.S. Pat. Nos. 4,485,045 and 4,544,545. Liposomes with enhanced circulation time are disclosed in U.S. Pat. No. 5,013,556.

Particularly useful liposomes can be generated by the reverse-phase evaporation method with a lipid composition comprising phosphatidylcholine, cholesterol, and PEG-derivatized phosphatidylethanolamine (PEG-PE). Liposomes are extruded through filters of defined pore size to yield liposomes with the desired diameter. Fab' fragments of the antibody of the present invention can be conjugated to the liposomes as described in Martin et al., *J. Biol. Chem.*, 257: 286-288 (1982) via a disulfide-interchange reaction. A chemotherapeutic agent (such as Doxorubicin) is optionally contained within the liposome. See Gabizon et al., *J. National Cancer Inst.*, 81(19): 1484 (1989).

9. Pharmaceutical Compositions of Antibodies

Antibodies specifically binding a PRO19598 polypeptide identified herein, as well as other molecules identified by the screening assays disclosed hereinbefore, can be administered for the treatment of various disorders in the form of pharmaceutical compositions.

If the PRO19598 polypeptide is intracellular and whole antibodies are used as inhibitors, internalizing antibodies are preferred. However, lipofections or liposomes can also be used to deliver the antibody, or an antibody fragment, into cells. Where antibody fragments are used, the smallest inhibitory fragment that specifically binds to the binding domain of the target protein is preferred. For example, based upon the variable-region sequences of an antibody, peptide molecules can be designed that retain the ability to bind the target protein sequence. Such peptides can be synthesized chemically and/or produced by recombinant DNA technology. See, e.g., Marasco et al., *Proc. Natl. Acad. Sci. USA*, 90: 7889-7893 (1993). The formulation herein may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each other. Alternatively, or in addition, the composition may comprise an agent that enhances its function, such as, for example, a cytotoxic agent, cytokine, chemotherapeutic agent, or growth-inhibitory agent. Such molecules are suitably present in combination in amounts that are effective for the purpose intended.

The active ingredients may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles, and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's *Pharmaceutical Sciences*, supra.

The formulations to be used for in vivo administration must be sterile. This is readily accomplished by filtration through sterile filtration membranes.

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semiper-

meable matrices of solid hydrophobic polymers containing the antibody, which matrices are in the form of shaped articles, e.g., films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 3,773,919), copolymers of L-glutamic acid and γ ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT™ (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods. When encapsulated antibodies remain in the body for a long time, they may denature or aggregate as a result of exposure to moisture at 37° C., resulting in a loss of biological activity and possible changes in immunogenicity. Rational strategies can be devised for stabilization depending on the mechanism involved. For example, if the aggregation mechanism is discovered to be intermolecular S—S bond formation through thio-disulfide interchange, stabilization may be achieved by modifying sulfhydryl residues, lyophilizing from acidic solutions, controlling moisture content, using appropriate additives, and developing specific polymer matrix compositions.

G. Uses for anti-PRO19598 Antibodies

The anti-PRO19598 antibodies of the invention have various utilities. For example, anti-PRO19598 antibodies may be used in diagnostic assays for PRO19598, e.g., detecting its expression (and in some cases, differential expression) in specific cells, tissues, or serum. Various diagnostic assay techniques known in the art may be used, such as competitive binding assays, direct or indirect sandwich assays and immunoprecipitation assays conducted in either heterogeneous or homogeneous phases [Zola, *Monoclonal Antibodies: A Manual of Techniques*, CRC Press, Inc. (1987) pp. 147-158]. The antibodies used in the diagnostic assays can be labeled with a detectable moiety. The detectable moiety should be capable of producing, either directly or indirectly, a detectable signal. For example, the detectable moiety may be a radioisotope, such as ^3H , ^{14}C , ^{32}P , ^{35}S , or ^{125}I , a fluorescent or chemiluminescent compound, such as fluorescein isothiocyanate, rhodamine, or luciferin, or an enzyme, such as alkaline phosphatase, beta-galactosidase or horseradish peroxidase. Any method known in the art for conjugating the antibody to the detectable moiety may be employed, including those methods described by Hunter et al., *Nature*, 144:945 (1962); David et al., *Biochemistry*, 13:1014 (1974); Pain et al., *J. Immunol. Meth.*, 40:219 (1981); and Nygren, *J. Histochem. and Cytochem.*, 30:407 (1982).

Anti-PRO19598 antibodies also are useful for the affinity purification of PRO19598 from recombinant cell culture or natural sources. In this process, the antibodies against PRO19598 are immobilized on a suitable support, such as Sephadex resin or filter paper, using methods well known in the art. The immobilized antibody then is contacted with a sample containing the PRO19598 to be purified, and thereafter the support is washed with a suitable solvent that will remove substantially all the material in the sample except the PRO19598, which is bound to the immobilized antibody. Finally, the support is washed with another suitable solvent that will release the PRO19598 from the antibody.

The following examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

All patent and literature references cited in the present specification are hereby incorporated by reference in their entirety.

EXAMPLES

Commercially available reagents referred to in the examples were used according to manufacturer's instruc-

reverse PCR primer

(SEQ ID NO:4)
>H3 ecd.snrl 5'TAAGTGCCTCCAGGTGCC 3'

Additionally, a synthetic oligonucleotide hybridization probe was constructed from the consensus DNA132879 sequence which had the following nucleotide sequence hybridization probe

5'TCCTCCAGTCATGAATATAACCCAAGTCAATGGCTCTTTGTTGGTAATTCTC 3' (SEQ ID NO:5)

tions unless otherwise indicated. The source of those cells identified in the following examples, and throughout the specification, by ATCC accession numbers is the American Type Culture Collection, Manassas, Va.

EXAMPLE 1

Isolation of cDNA Clones Encoding a Human PRO19598

The extracellular domain (ECD) sequences (including the secretion signal sequence, if any) from about 950 known secreted proteins from the Swiss-Prot public database were used to search sequence databases. The databases included public databases (e.g., GenBank) In this instance, genomic DNA sequence from GenBank was analyzed using the gene prediction program GENSCAN, licenced from Stanford University. GENSCAN analysis predicts gene coding regions, creating sequences which can be subjected to the ECD search. The search was performed using the computer program BLAST or BLAST2 [Altschul et al., *Methods in Enzymology*, 266:460-480 (1996)] as a comparison of the ECD protein sequences to a 6 frame translation of the sequences. Those comparisons resulting in a BLAST score of 70 (or in some cases, 90) or greater that did not encode known proteins were clustered and assembled into consensus DNA sequences with the program "phrap" (Phil Green, University of Washington, Seattle, Wash.).

A consensus DNA sequence was assembled. This consensus sequence is herein designated DNA132879.

Based on the DNA132879 consensus sequence, oligonucleotides were synthesized: 1) to identify by PCR a cDNA library that contained the sequence of interest, and 2) for use as probes to isolate a clone of the full-length coding sequence for PRO19598. Forward and reverse PCR primers generally range from 20 to 30 nucleotides and are often designed to give a PCR product of about 100-1000 bp in length. The probe sequences are typically 40-55 bp in length. In some cases, additional oligonucleotides are synthesized when the consensus sequence is greater than about 1-1.5 kbp. In order to screen several libraries for a full-length clone, DNA from the libraries was screened by PCR amplification, as per Ausubel et al., *Current Protocols in Molecular Biology*, supra, with the PCR primer pair. A positive library was then used to isolate clones encoding the gene of interest using the probe oligonucleotide and one of the primer pairs.

PCR primers (forward and reverse) were synthesized:
forward PCR primer

(SEQ ID NO:3)
>H3 ecd.snrl 5'CTGGCAACAGCAGTGTCTATTTGTGC 3'

A pool of 50 different human cDNA libraries from various tissues was used in cloning. The cDNA libraries used to isolate the cDNA clones were constructed by standard methods using commercially available reagents such as those from Invitrogen, San Diego, Calif. The cDNA was primed with oligo dT containing a NotI site, linked with blunt to SalI hemikinased adaptors, cleaved with NotI, sized appropriately by gel electrophoresis, and cloned in a defined orientation into a suitable cloning vector (such as pRKB or pRKD; pRK5B is a precursor of pRK5D that does not contain the SfiI site; see, Holmes et al., *Science*, 253:1278-1280 (1991)) in the unique XhoI and NotI sites.

DNA sequencing of the clones isolated as described above gave the full-length DNA sequence for a full-length PRO19598 polypeptide (designated herein as DNA145887-2849 [FIG. 1, SEQ ID NO: 1) and the derived protein sequence for that PRO19598 polypeptide.

The full length clone identified above contained a single open reading frame with an apparent translational initiation site at nucleotide positions 241-243 and a stop signal at nucleotide positions 1027-1030 (FIG. 1, SEQ ID NO:1). The predicted polypeptide precursor is 262 amino acids long, has a calculated molecular weight of approximately 30419 daltons and an estimated pI of approximately 8.44. Analysis of the full-length PRO19598 sequence shown in FIG. 2 (SEQ ID NO:2) evidences the presence of a variety of important polypeptide domains as shown in FIG. 2, wherein the locations given for those important polypeptide domains are approximate as described above. Clone DNA145887-2849 has been deposited with ATCC on Mar. 21, 2000 and is assigned ATCC deposit no. PTA-1532.

An analysis of the Dayhoff database (version 35.45 SwissProt 35), using the ALIGN-2 sequence alignment analysis of the full-length sequence shown in FIG. 2 (SEQ ID NO:2), evidenced sequence identity between the PRO19598 amino acid sequence and the following Dayhoff sequences:AF184971_1.

EXAMPLE 2

Use of PRO19598 as a Hybridization Probe

The following method describes use of a nucleotide sequence encoding PRO19598 as a hybridization probe.

DNA comprising the coding sequence of full-length or mature PRO19598 as disclosed herein is employed as a probe to screen for homologous DNAs (such as those encoding naturally-occurring variants of PRO19598) in human tissue cDNA libraries or human tissue genomic libraries.

Hybridization and washing of filters containing either library DNAs is performed under the following high stringency conditions. Hybridization of radiolabeled PRO19598-derived probe to the filters is performed in a solution of 50% formamide, 5×SSC, 0.1% SDS, 0.1% sodium pyrophosphate, 50 mM sodium phosphate, pH 6.8,

2xDenhardt's solution, and 10% dextran sulfate at 42° C. for 20 hours. Washing of the filters is performed in an aqueous solution of 0.1xSSC and 0.1% SDS at 42° C.

DNAs having a desired sequence identity with the DNA encoding full-length native sequence PRO19598 can then be identified using standard techniques known in the art.

EXAMPLE 3

Expression of PRO19598 in *E. coli*

This example illustrates preparation of an unglycosylated form of PRO19598 by recombinant expression in *E. coli*.

The DNA sequence encoding PRO19598 is initially amplified using selected PCR primers. The primers should contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector. A variety of expression vectors may be employed. An example of a suitable vector is pBR322 (derived from *E. coli*; see Bolivar et al., *Gene*, 2:95 (1977)) which contains genes for ampicillin and tetracycline resistance. The vector is digested with restriction enzyme and dephosphorylated. The PCR amplified sequences are then ligated into the vector. The vector will preferably include sequences which encode for an antibiotic resistance gene, a trp promoter, a polyhis leader (including the first six STII codons, polyhis sequence, and enterokinase cleavage site), the PRO19598 coding region, lambda transcriptional terminator, and an argU gene.

The ligation mixture is then used to transform a selected *E. coli* strain using the methods described in Sambrook et al., supra. Transformants are identified by their ability to grow on LB plates and antibiotic resistant colonies are then selected. Plasmid DNA can be isolated and confirmed by restriction analysis and DNA sequencing.

Selected clones can be grown overnight in liquid culture medium such as LB broth supplemented with antibiotics. The overnight culture may subsequently be used to inoculate a larger scale culture. The cells are then grown to a desired optical density, during which the expression promoter is turned on.

After culturing the cells for several more hours, the cells can be harvested by centrifugation. The cell pellet obtained by the centrifugation can be solubilized using various agents known in the art, and the solubilized PRO19598 protein can then be purified using a metal chelating column under conditions that allow tight binding of the protein.

PRO19598 may be expressed in *E. coli* in a poly-His tagged form, using the following procedure. The DNA encoding PRO19598 is initially amplified using selected PCR primers. The primers will contain restriction enzyme sites which correspond to the restriction enzyme sites on the selected expression vector, and other useful sequences providing for efficient and reliable translation initiation, rapid purification on a metal chelation column, and proteolytic removal with enterokinase. The PCR-amplified, poly-His tagged sequences are then ligated into an expression vector, which is used to transform an *E. coli* host based on strain 52 (W3110 fuhA(tonA) lon galE rpoHts(htpRts) clpP(lacIq). Transformants are first grown in LB containing 50 mg/ml carbenicillin at 30° C. with shaking until an O.D.600 of 3–5 is reached. Cultures are then diluted 50–100 fold into CRAP media (prepared by mixing 3.57 g (NH₄)₂SO₄, 0.71 g sodium citrate.2H₂O, 1.07 g KCl, 5.36 g Difco yeast extract, 5.36 g Sheffield bycase SF in 500 mL water, as well as 110 mM MPOS, pH 7.3, 0.55% (w/v) glucose and 7 mM MgSO₄) and grown for approximately 20–30 hours at 30° C. with shaking. Samples are removed to verify expression by SDS-PAGE analysis, and the bulk culture is centrifuged to pellet the cells. Cell pellets are frozen until purification and refolding.

E. coli paste from 0.5 to 1 L fermentations (6–10 g pellets) is resuspended in 10 volumes (w/v) in 7 M guanidine, 20 mM Tris, pH 8 buffer. Solid sodium sulfite and sodium tetrathionate is added to make final concentrations of 0.1M and 0.02 M, respectively, and the solution is stirred overnight at 4° C. This step results in a denatured protein with all cysteine residues blocked by sulfitolization. The solution is centrifuged at 40,000 rpm in a Beckman Ultracentrifuge for 30 min. The supernatant is diluted with 3–5 volumes of metal chelate column buffer (6 M guanidine, 20 mM Tris, pH 7.4) and filtered through 0.22 micron filters to clarify. The clarified extract is loaded onto a 5 ml Qiagen Ni-NTA metal chelate column equilibrated in the metal chelate column buffer. The column is washed with additional buffer containing 50 mM imidazole (Calbiochem, Utrol grade), pH 7.4. The protein is eluted with buffer containing 250 mM imidazole. Fractions containing the desired protein are pooled and stored at 4° C. Protein concentration is estimated by its absorbance at 280 nm using the calculated extinction coefficient based on its amino acid sequence.

The proteins are refolded by diluting the sample slowly into freshly prepared refolding buffer consisting of: 20 mM Tris, pH 8.6, 0.3 M NaCl, 2.5 M urea, 5 mM cysteine, 20 mM glycine and 1 mM EDTA. Refolding volumes are chosen so that the final protein concentration is between 50 to 100 micrograms/ml. The refolding solution is stirred gently at 4° C. for 12–36 hours. The refolding reaction is quenched by the addition of TFA to a final concentration of 0.4% (pH of approximately 3). Before further purification of the protein, the solution is filtered through a 0.22 micron filter and acetonitrile is added to 2–10% final concentration. The refolded protein is chromatographed on a Poros R1/H reversed phase column using a mobile buffer of 0.1% TFA with elution with a gradient of acetonitrile from 10 to 80%. Aliquots of fractions with A280 absorbance are analyzed on SDS polyacrylamide gels and fractions containing homogeneous refolded protein are pooled. Generally, the properly refolded species of most proteins are eluted at the lowest concentrations of acetonitrile since those species are the most compact with their hydrophobic interiors shielded from interaction with the reversed phase resin. Aggregated species are usually eluted at higher acetonitrile concentrations. In addition to resolving misfolded forms of proteins from the desired form, the reversed phase step also removes endotoxin from the samples.

Fractions containing the desired folded PRO19598 polypeptide are pooled and the acetonitrile removed using a gentle stream of nitrogen directed at the solution. Proteins are formulated into 20 mM Hepes, pH 6.8 with 0.14 M sodium chloride and 4% mannitol by dialysis or by gel filtration using G25 Superfine (Pharmacia) resins equilibrated in the formulation buffer and sterile filtered.

PRO19598 polypeptides disclosed herein were successfully expressed as described above.

EXAMPLE 4

Expression of PRO19598 in Mammalian Cells

This example illustrates preparation of a potentially glycosylated form of PRO19598 by recombinant expression in mammalian cells.

The vector, pRK5 (see EP 307,247, published Mar. 15, 1989), is employed as the expression vector. Optionally, the PRO19598 DNA is ligated into pRK5 with selected restriction enzymes to allow insertion of the PRO19598 DNA using ligation methods such as described in Sambrook et al., supra. The resulting vector is called pRK5-PRO19598.

In one embodiment, the selected host cells may be 293 cells. Human 293 cells (ATCC CCL 1573) are grown to

confluence in tissue culture plates in medium such as DMEM supplemented with fetal calf serum and optionally, nutrient components and/or antibiotics. About 10 μ g pRK5-PRO19598 DNA is mixed with about 1 μ g DNA encoding the VARNA gene [Thimmappaya et al., *Cell*, 31:543 (1982)] and dissolved in 500 μ l of 1 mM Tris-HCl, 0.1 mM EDTA, 0.227 M CaCl₂. To this mixture is added, dropwise, 500 μ l of 50 mM HEPES (pH 7.35), 280 mM NaCl, 1.5 mM NaPO₄, and a precipitate is allowed to form for 10 minutes at 25° C. The precipitate is suspended and added to the 293 cells and allowed to settle for about four hours at 37° C. The culture medium is aspirated off and 2 ml of 20% glycerol in PBS is added for 30 seconds. The 293 cells are then washed with serum free medium, fresh medium is added and the cells are incubated for about 5 days.

Approximately 24 hours after the transfections, the culture medium is removed and replaced with culture medium (alone) or culture medium containing 200 μ Ci/ml ³⁵S-cysteine and 200 μ Ci/ml ³⁵S-methionine. After a 12 hour incubation, the conditioned medium is collected, concentrated on a spin filter, and loaded onto a 15% SDS gel. The processed gel may be dried and exposed to film for a selected period of time to reveal the presence of PRO19598 polypeptide. The cultures containing transfected cells may undergo further incubation (in serum free medium) and the medium is tested in selected bioassays.

In an alternative technique, PRO19598 may be introduced into 293 cells transiently using the dextran sulfate method described by Somparyrac et al., *Proc. Natl. Acad. Sci.*, 12:7575 (1981). 293 cells are grown to maximal density in a spinner flask and 700 μ g pRK5-PRO19598 DNA is added. The cells are first concentrated from the spinner flask by centrifugation and washed with PBS. The DNA-dextran precipitate is incubated on the cell pellet for four hours. The cells are treated with 20% glycerol for 90 seconds, washed with tissue culture medium, and re-introduced into the spinner flask containing tissue culture medium, 5 μ g/ml bovine insulin and 0.1 μ g/ml bovine transferrin. After about four days, the conditioned media is centrifuged and filtered to remove cells and debris. The sample containing expressed PRO19598 can then be concentrated and purified by any selected method, such as dialysis and/or column chromatography.

In another embodiment, PRO19598 can be expressed in CHO cells. The pRK5-PRO19598 can be transfected into CHO cells using known reagents such as CaPO₄ or DEAE-dextran. As described above, the cell cultures can be incubated, and the medium replaced with culture medium (alone) or medium containing a radiolabel such as ³⁵S-methionine. After determining the presence of PRO19598 polypeptide, the culture medium may be replaced with serum free medium. Preferably, the cultures are incubated for about 6 days, and then the conditioned medium is harvested. The medium containing the expressed PRO19598 can then be concentrated and purified by any selected method.

Epitope-tagged PRO19598 may also be expressed in host CHO cells. The PRO19598 may be subcloned out of the pRK5 vector. The subclone insert can undergo PCR to fuse in frame with a selected epitope tag such as a poly-his tag into a Baculovirus expression vector. The poly-his tagged PRO19598 insert can then be subcloned into a SV40 driven vector containing a selection marker such as DHFR for selection of stable clones. Finally, the CHO cells can be transfected (as described above) with the SV40 driven vector. Labeling may be performed, as described above, to verify expression. The culture medium containing the

expressed poly-His tagged PRO19598 can then be concentrated and purified by any selected method, such as by Ni²⁺-chelate affinity chromatography.

PRO19598 may also be expressed in CHO and/or COS cells by a transient expression procedure or in CHO cells by another stable expression procedure.

Stable expression in CHO cells is performed using the following procedure. The proteins are expressed as an IgG construct (immunoadhesin), in which the coding sequences for the soluble forms (e.g. extracellular domains) of the respective proteins are fused to an IgG1 constant region sequence containing the hinge, CH2 and CH2 domains and/or is a poly-His tagged form.

Following PCR amplification, the respective DNAs are subcloned in a CHO expression vector using standard techniques as described in Ausubel et al., *Current Protocols of Molecular Biology*, Unit 3.16, John Wiley and Sons (1997). CHO expression vectors are constructed to have compatible restriction sites 5' and 3' of the DNA of interest to allow the convenient shuttling of cDNA's. The vector used expression in CHO cells is as described in Lucas et al., *Nucl. Acids Res.* 24:9 (1774-1779 (1996), and uses the SV40 early promoter/enhancer to drive expression of the cDNA of interest and dihydrofolate reductase (DHFR). DHFR expression permits selection for stable maintenance of the plasmid following transfection.

Twelve micrograms of the desired plasmid DNA is introduced into approximately 10 million CHO cells using commercially available transfection reagents Superfect® (Qiagen), Dosper® or Eugene® (Boehringer Mannheim). The cells are grown as described in Lucas et al., *supra*. Approximately 3×10⁷ cells are frozen in an ampule for further growth and production as described below.

The ampules containing the plasmid DNA are thawed by placement into water bath and mixed by vortexing. The contents are pipetted into a centrifuge tube containing 10 mL of media and centrifuged at 1000 rpm for 5 minutes. The supernatant is aspirated and the cells are resuspended in 10 mL of selective media (0.2 μ m filtered PS20 with 5% 0.2 μ m diafiltered fetal bovine serum). The cells are then aliquoted into a 100 mL spinner containing 90 mL of selective media. After 1-2 days, the cells are transferred into a 250 mL spinner filled with 150 mL selective growth medium and incubated at 37° C. After another 2-3 days, 250 mL, 500 mL and 2000 mL spinners are seeded with 3×10⁵ cells/mL. The cell media is exchanged with fresh media by centrifugation and resuspension in production medium. Although any suitable CHO media may be employed, a production medium described in U.S. Pat. No. 5,122,469, issued Jun. 16, 1992 may actually be used. A 3L production spinner is seeded at 1.2×10⁶ cells/mL. On day 0, the cell number pH is determined. On day 1, the spinner is sampled and sparging with filtered air is commenced. On day 2, the spinner is sampled, the temperature shifted to 33° C., and 30 mL of 500 g/L glucose and 0.6 mL of 10% antifoam (e.g., 35% polydimethylsiloxane em Dow Corning 365 Medical Grade Emulsion) taken. Throughout the production, the pH is adjusted as necessary to keep it at around 7.2. After 10 days, or until the viability dropped below 70%, the cell culture is harvested by centrifugation and filtering through a 0.22 μ m filter. The filtrate was either stored at 4° C. or immediately loaded onto columns for purification.

For the poly-His tagged constructs, the proteins are purified using a Ni-NTA column (Qiagen). Before purification, imidazole is added to the conditioned media to a concentration of 5 mM. The conditioned media is pumped onto a 6 ml Ni-NTA column equilibrated in 20 mM Hepes, pH 7.4,

buffer containing 0.3 M NaCl and 5 mM imidazole at a flow rate of 4–5 ml/min. at 4° C. After loading, the column is washed with additional equilibration buffer and the protein eluted with equilibration buffer containing 0.25 M imidazole. The highly purified protein is subsequently desalted into a storage buffer containing 10 mM Hepes, 0.14 M NaCl and 4% mannitol, pH 6.8, with a 25 ml G25 Superfine (Pharmacia) column and stored at –80° C.

Immunoadhesin (Fc-containing) constructs are purified from the conditioned media as follows. The conditioned medium is pumped onto a 5 ml Protein A column (Pharmacia) which had been equilibrated in 20 mM Na phosphate buffer, pH 6.8. After loading, the column is washed extensively with equilibration buffer before elution with 100 mM citric acid, pH 3.5. The eluted protein is immediately neutralized by collecting 1 ml fractions into tubes containing 275 μ L of 1 M Tris buffer, pH 9. The highly purified protein is subsequently desalted into storage buffer as described above for the poly-His tagged proteins. The homogeneity is assessed by SDS polyacrylamide gels and by N-terminal amino acid sequencing by Edman degradation.

EXAMPLE 5

Expression of PRO19598 in Yeast

The following method describes recombinant expression of PRO19598 in yeast.

First, yeast expression vectors are constructed for intracellular production or secretion of PRO19598 from the ADH2/GAPDH promoter. DNA encoding PRO19598 and the promoter is inserted into suitable restriction enzyme sites in the selected plasmid to direct intracellular expression of PRO19598. For secretion, DNA encoding PRO19598 can be cloned into the selected plasmid, together with DNA encoding the ADH2/GAPDH promoter, a native PRO19598 signal peptide or other mammalian signal peptide, or, for example, a yeast α -factor or invertase secretory signal/leader sequence, and linker sequences (if needed) for expression of PRO19598.

Yeast cells, such as yeast strain AB110, can then be transformed with the expression plasmids described above and cultured in selected fermentation media. The transformed yeast supernatants can be analyzed by precipitation with 10% trichloroacetic acid and separation by SDS-PAGE, followed by staining of the gels with Coomassie Blue stain.

Recombinant PRO19598 can subsequently be isolated and purified by removing the yeast cells from the fermentation medium by centrifugation and then concentrating the medium using selected cartridge filters. The concentrate containing PRO19598 may further be purified using selected column chromatography resins.

EXAMPLE 6

Expression of PRO19598 in Baculovirus-infected Insect Cells

The following method describes recombinant expression of PRO19598 in Baculovirus-infected insect cells.

The sequence coding for PRO19598 is fused upstream of an epitope tag contained within a baculovirus expression vector. Such epitope tags include poly-his tags and immunoglobulin tags (like Fc regions of IgG). A variety of plasmids may be employed, including plasmids derived from commercially available plasmids such as pVL1393 (Novagen). Briefly, the sequence encoding PRO19598 or the desired portion of the coding sequence of PRO19598 such as the sequence encoding the extracellular domain of a transmembrane protein or the sequence encoding the mature protein if the protein is extracellular is amplified by PCR with primers complementary to the 5' and 3' regions. The 5'

primer may incorporate flanking (selected) restriction enzyme sites. The product is then digested with those selected restriction enzymes and subcloned into the expression vector.

Recombinant baculovirus is generated by co-transfecting the above plasmid and BaculoGold™ virus DNA (Pharmingen) into *Spodoptera frugiperda* ("Sf9") cells (ATCC CRL 1711) using lipofectin (commercially available from GIBCO-BRL). After 4–5 days of incubation at 28° C., the released viruses are harvested and used for further amplifications. Viral infection and protein expression are performed as described by O'Reilly et al., *Baculovirus expression vectors: A Laboratory Manual*, Oxford: Oxford University Press (1994).

Expressed poly-his tagged PRO19598 can then be purified, for example, by Ni^{2+} -chelate affinity chromatography as follows. Extracts are prepared from recombinant virus-infected Sf9 cells as described by Rupert et al., *Nature*, 362:175–179 (1993). Briefly, Sf9 cells are washed, resuspended in sonication buffer (25 mM Hepes, pH 7.9; 12.5 mM MgCl_2 ; 0.1 mM EDTA; 10% glycerol; 0.1% NP-40; 0.4 M KCl), and sonicated twice for 20 seconds on ice. The sonicates are cleared by centrifugation, and the supernatant is diluted 50-fold in loading buffer (50 mM phosphate, 300 mM NaCl, 10% glycerol, pH 7.8) and filtered through a 0.45 μ m filter. A Ni^{2+} -NTA agarose column (commercially available from Qiagen) is prepared with a bed volume of 5 mL, washed with 25 mL of water and equilibrated with 25 mL of loading buffer. The filtered cell extract is loaded onto the column at 0.5 mL per minute. The column is washed to baseline A_{280} with loading buffer, at which point fraction collection is started. Next, the column is washed with a secondary wash buffer (50 mM phosphate; 300 mM NaCl, 10% glycerol, pH 6.0), which elutes nonspecifically bound protein. After reaching A_{280} baseline again, the column is developed with a 0 to 500 mM Imidazole gradient in the secondary wash buffer. One mL fractions are collected and analyzed by SDS-PAGE and silver staining or Western blot with Ni^{2+} -NTA-conjugated to alkaline phosphatase (Qiagen). Fractions containing the eluted His_{10} -tagged PRO19598 are pooled and dialyzed against loading buffer.

Alternatively, purification of the IgG tagged (or Fc tagged) PRO19598 can be performed using known chromatography techniques, including for instance, Protein A or protein G column chromatography.

PRO19598 polypeptides disclosed herein were successfully expressed as described above.

EXAMPLE 7

Preparation of Antibodies that Bind PRO19598

This example illustrates preparation of monoclonal antibodies which can specifically bind PRO19598.

Techniques for producing the monoclonal antibodies are known in the art and are described, for instance, in Goding, supra. Immunogens that may be employed include purified PRO 19598, fusion proteins containing PRO19598, and cells expressing recombinant PRO19598 on the cell surface. Selection of the immunogen can be made by the skilled artisan without undue experimentation.

Mice, such as Balb/c, are immunized with the PRO19598 immunogen emulsified in complete Freund's adjuvant and injected subcutaneously or intraperitoneally in an amount from 1–100 micrograms. Alternatively, the immunogen is emulsified in MPL-TDM adjuvant (Ribi Immunochemical Research, Hamilton, Mont.) and injected into the animal's hind foot pads. The immunized mice are then boosted 10 to 12 days later with additional immunogen emulsified in the selected adjuvant. Thereafter, for several weeks, the mice

may also be boosted with additional immunization injections. Serum samples may be periodically obtained from the mice by retro-orbital bleeding for testing in ELISA assays to detect anti-PRO19598 antibodies.

After a suitable antibody titer has been detected, the animals "positive" for antibodies can be injected with a final intravenous injection of PRO19598. Three to four days later, the mice are sacrificed and the spleen cells are harvested. The spleen cells are then fused (using 35% polyethylene glycol) to a selected murine myeloma cell line such as P3X63AgU.1, available from ATCC, No. CRL 1597. The fusions generate hybridoma cells which can then be plated in 96 well tissue culture plates containing HAT (hypoxanthine, aminopterin, and thymidine) medium to inhibit proliferation of non-fused cells, myeloma hybrids, and spleen cell hybrids.

The hybridoma cells will be screened in an ELISA for reactivity against PRO 19598. Determination of "positive" hybridoma cells secreting the desired monoclonal antibodies against PRO19598 is within the skill in the art.

The positive hybridoma cells can be injected intraperitoneally into syngeneic Balb/c mice to produce ascites containing the anti-PRO19598 monoclonal antibodies. Alternatively, the hybridoma cells can be grown in tissue culture flasks or roller bottles. Purification of the monoclonal antibodies produced in the ascites can be accomplished using ammonium sulfate precipitation, followed by gel exclusion chromatography. Alternatively, affinity chromatography based upon binding of antibody to protein A or protein G can be employed.

EXAMPLE 8

Purification of PRO19598 Polypeptides Using Specific Antibodies

Native or recombinant PRO19598 polypeptides may be purified by a variety of standard techniques in the art of protein purification. For example, pro-PRO19598 polypeptide, mature PRO19598 polypeptide, or pre-PRO19598 polypeptide is purified by immunoaffinity chromatography using antibodies specific for the PRO19598 polypeptide of interest. In general, an immunoaffinity column is constructed by covalently coupling the anti-PRO19598 polypeptide antibody to an activated chromatographic resin.

Polyclonal immunoglobulins are prepared from immune sera either by precipitation with ammonium sulfate or by purification on immobilized Protein A (Pharmacia LKB Biotechnology, Piscataway, N.J.). Likewise, monoclonal antibodies are prepared from mouse ascites fluid by ammonium sulfate precipitation or chromatography on immobilized Protein A. Partially purified immunoglobulin is covalently attached to a chromatographic resin such as CnBr-activated SEPHAROSE™ (Pharmacia LKB Biotechnology). The antibody is coupled to the resin, the resin is blocked, and the derivative resin is washed according to the manufacturer's instructions.

Such an immunoaffinity column is utilized in the purification of PRO19598 polypeptide by preparing a fraction from cells containing PRO19598 polypeptide in a soluble form. This preparation is derived by solubilization of the whole cell or of a subcellular fraction obtained via differential centrifugation by the addition of detergent or by other methods well known in the art. Alternatively, soluble PRO19598 polypeptide containing a signal sequence may be secreted in useful quantity into the medium in which the cells are grown.

A soluble PRO19598 polypeptide-containing preparation is passed over the immunoaffinity column, and the column

is washed under conditions that allow the preferential absorbance of PRO19598 polypeptide (e.g., high ionic strength buffers in the presence of detergent). Then, the column is eluted under conditions that disrupt antibody/PRO19598 polypeptide binding (e.g., a low pH buffer such as approximately pH 2-3, or a high concentration of a chaotrope such as urea or thiocyanate ion), and PRO19598 polypeptide is collected.

EXAMPLE 9

Drug Screening

This invention is particularly useful for screening compounds by using PRO19598 polypeptides or binding fragment thereof in any of a variety of drug screening techniques. The PRO19598 polypeptide or fragment employed in such a test may either be free in solution, affixed to a solid support, borne on a cell surface, or located intracellularly. One method of drug screening utilizes eukaryotic or prokaryotic host cells which are stably transformed with recombinant nucleic acids expressing the PRO19598 polypeptide or fragment. Drugs are screened against such transformed cells in competitive binding assays. Such cells, either in viable or fixed form, can be used for standard binding assays. One may measure, for example, the formation of complexes between PRO19598 polypeptide or a fragment and the agent being tested. Alternatively, one can examine the diminution in complex formation between the PRO19598 polypeptide and its target cell or target receptors caused by the agent being tested.

Thus, the present invention provides methods of screening for drugs or any other agents which can affect a PRO19598 polypeptide-associated disease or disorder. These methods comprise contacting such an agent with an PRO19598 polypeptide or fragment thereof and assaying (i) for the presence of a complex between the agent and the PRO19598 polypeptide or fragment, or (ii) for the presence of a complex between the PRO19598 polypeptide or fragment and the cell, by methods well known in the art. In such competitive binding assays, the PRO19598 polypeptide or fragment is typically labeled. After suitable incubation, free PRO19598 polypeptide or fragment is separated from that present in bound form, and the amount of free or uncomplexed label is a measure of the ability of the particular agent to bind to PRO19598 polypeptide or to interfere with the PRO19598 polypeptide/cell complex.

Another technique for drug screening provides high throughput screening for compounds having suitable binding affinity to a polypeptide and is described in detail in WO 84/03564, published on Sep. 13, 1984. Briefly stated, large numbers of different small peptide test compounds are synthesized on a solid substrate, such as plastic pins or some other surface. As applied to a PRO19598 polypeptide, the peptide test compounds are reacted with PRO19598 polypeptide and washed. Bound PRO19598 polypeptide is detected by methods well known in the art. Purified PRO19598 polypeptide can also be coated directly onto plates for use in the aforementioned drug screening techniques. In addition, non-neutralizing antibodies can be used to capture the peptide and immobilize it on the solid support.

This invention also contemplates the use of competitive drug screening assays in which neutralizing antibodies capable of binding PRO19598 polypeptide specifically compete with a test compound for binding to PRO19598 polypeptide or fragments thereof. In this manner, the antibodies can be used to detect the presence of any peptide which shares one or more antigenic determinants with PRO19598 polypeptide.

EXAMPLE 10

Rational Drug Design

The goal of rational drug design is to produce structural analogs of biologically active polypeptide of interest (i.e., a PRO19598 polypeptide) or of small molecules with which they interact, e.g., agonists, antagonists, or inhibitors. Any of these examples can be used to fashion drugs which are more active or stable forms of the PRO19598 polypeptide or which enhance or interfere with the function of the PRO19598 polypeptide in vivo (cf., Hodgson, *Bio/Technology*, 9: 19-21(1991)).

In one approach, the three-dimensional structure of the PRO19598 polypeptide, or of an PRO19598 polypeptide-inhibitor complex, is determined by x-ray crystallography, by computer modeling or, most typically, by a combination of the two approaches. Both the shape and charges of the PRO19598 polypeptide must be ascertained to elucidate the structure and to determine active site(s) of the molecule. Less often, useful information regarding the structure of the PRO19598 polypeptide may be gained by modeling based on the structure of homologous proteins. In both cases, relevant structural information is used to design analogous PRO19598 polypeptide-like molecules or to identify efficient inhibitors. Useful examples of rational drug design may include molecules which have improved activity or stability as shown by Braxton and Wells, *Biochemistry*, 31:7796-7801 (1992) or which act as inhibitors, agonists, or antagonists of native peptides as shown by Athauda et al., *J. Biochem.*, 113:742-746 (1993).

It is also possible to isolate a target-specific antibody, selected by functional assay, as described above, and then to solve its crystal structure. This approach, in principle, yields a pharmacore upon which subsequent drug design can be based. It is possible to bypass protein crystallography altogether by generating anti-idiotypic antibodies (anti-ids) to a functional, pharmacologically active antibody. As a mirror image of a mirror image, the binding site of the anti-ids would be expected to be an analog of the original receptor. The anti-id could then be used to identify and isolate peptides from banks of chemically or biologically produced peptides. The isolated peptides would then act as the pharmacore.

By virtue of the present invention, sufficient amounts of the PRO19598 polypeptide may be made available to perform such analytical studies as X-ray crystallography. In addition, knowledge of the PRO19598 polypeptide amino acid sequence provided herein will provide guidance to those employing computer modeling techniques in place of or in addition to x-ray crystallography.

EXAMPLE 11

Tissue Expression Distribution

Oligonucleotide probes were constructed from the PRO19598 polypeptide-encoding nucleotide sequence shown in FIG. 1 for use in quantitative PCR amplification reactions. The oligonucleotide probes were chosen so as to give an approximately 200-600 base pair amplified fragment from the 3' end of its associated template in a standard PCR reaction. The oligonucleotide probes were employed in standard quantitative PCR amplification reactions with cDNA libraries isolated from different human adult and/or fetal tissue sources and analyzed by agarose gel electrophoresis so as to obtain a quantitative determination of the level of expression of the PRO19598 polypeptide-encoding nucleic acid in the various tissues tested. Knowledge of the expression pattern or the differential expression of the PRO19598 polypeptide-encoding nucleic acid in various different human tissue types provides a diagnostic marker

useful for tissue typing, with or without other tissue-specific markers, for determining the primary tissue source of a metastatic tumor, disease diagnosis, and the like. These assays provided the following results:

DNA Molecule	
Tissues w/Significant Expression	
DNA145887-2849	Highly expressed in cartilage, placenta and spleen. It is expressed in testis, adrenal gland and prostate.
Tissues w/o Significant Expression	
DNA145887-2849	Is not expressed in HUVEC, colon tumor, heart, bone marrow, aortic endothelial cells or uterus.

EXAMPLE 12

Identification of Receptor/Ligand Interactions

In this assay, various PRO polypeptides are tested for ability to bind to a panel of potential receptor molecules for the purpose of identifying receptor/ligand interactions. The identification of a ligand for a known receptor, a receptor for a known ligand or a novel receptor/ligand pair is useful for a variety of indications including, for example, targeting bioactive molecules (linked to the ligand or receptor) to a cell known to express the receptor or ligand, use of the receptor or ligand as a reagent to detect the presence of the ligand or receptor in a composition suspected of containing the same, wherein the composition may comprise cells suspected of expressing the ligand or receptor, modulating the growth of or another biological or immunological activity of a cell known to express or respond to the receptor or ligand, modulating the immune response of cells or toward cells that express the receptor or ligand, allowing the preparation of agonists, antagonists and/or antibodies directed against the receptor or ligand which will modulate the growth of or a biological or immunological activity of a cell expressing the receptor or ligand, and various other indications which will be readily apparent to the ordinarily skilled artisan.

The assay is performed as follows. A PRO polypeptide of the present invention suspected of being a ligand for a receptor is expressed as a fusion protein containing the Fc domain of human IgG (an Immunoadhesin). Receptor-ligand binding is detected by allowing interaction of the immunoadhesin polypeptide with cells (e.g. Cos cells) expressing candidate PRO polypeptide receptors and visualization of bound immunoadhesin with fluorescent reagents directed toward the Fc fusion domain and examination by microscope. Cells expressing candidate receptors are produced by transient transfection, in parallel, of defined subsets of a library of cDNA expression vectors encoding PRO polypeptides that may function as receptor molecules. Cells are then incubated for 1 hour in the presence of the PRO polypeptide immunoadhesin being tested for possible receptor binding. The cells are then washed and fixed with paraformaldehyde. The cells are then incubated with fluorescent conjugated antibody directed against the Fc portion of the PRO polypeptide immunoadhesin (e.g. FITC conjugated goat anti-human-Fc antibody). The cells are then washed again and examined by microscope. A positive interaction is judged by the presence of fluorescent labeling of cells transfected with cDNA encoding a particular PRO polypeptide receptor or pool of receptors and an absence of similar fluorescent labeling of similarly prepared cells that have been transfected with other cDNA or pools of cDNA.

If a defined pool of cDNA expression vectors is judged to be positive for interaction with a PRO polypeptide immunoadhesin, the individual cDNA species that comprise the pool are tested individually (the pool is "broken down") to determine the specific cDNA that encodes a receptor able to interact with the PRO polypeptide immunoadhesin.

In another embodiment of this assay, an epitope-tagged potential ligand PRO polypeptide (e.g. 8 histidine "His" tag) is allowed to interact with a panel of potential receptor PRO polypeptide molecules that have been expressed as fusions with the Fc domain of human IgG (immunoadhesins). Following a 1 hour co-incubation with the epitope tagged PRO polypeptide, the candidate receptors are each immunoprecipitated with protein A beads and the beads are washed. Potential ligand interaction is determined by western blot analysis of the immunoprecipitated complexes with antibody directed towards the epitope tag. An interaction is judged to occur if a band of the anticipated molecular weight of the epitope tagged protein is observed in the western blot analysis with a candidate receptor, but is not observed to occur with the other members of the panel of potential receptors.

(1) PRO19598 binds to PRO3301 and vice versa.

EXAMPLE 13

Microarray Analysis to Detect Overexpression of PRO Polypeptides in Cancerous Tumors

Nucleic acid microarrays, often containing thousands of gene sequences, are useful for identifying differentially expressed genes in diseased tissues as compared to their normal counterparts. Using nucleic acid microarrays, test and control mRNA samples from test and control tissue samples are reverse transcribed and labeled to generate cDNA probes. The cDNA probes are then hybridized to an array of nucleic acids immobilized on a solid support. The array is configured such that the sequence and position of each member of the array is known. For example, a selection of genes known to be expressed in certain disease states may be arrayed on a solid support. Hybridization of a labeled probe with a particular array member indicates that the sample from which the probe was derived expresses that gene. If the hybridization signal of a probe from a test (disease tissue) sample is greater than hybridization signal of a probe from a control (normal tissue) sample, the gene or genes overexpressed in the disease tissue are identified. The implication of this result is that an overexpressed protein in a diseased tissue is useful not only as a diagnostic marker for the presence of the disease condition, but also as a therapeutic target for treatment of the disease condition.

The methodology of hybridization of nucleic acids and microarray technology is well known in the art. In the present example, the specific preparation of nucleic acids for hybridization and probes, slides, and hybridization conditions are all detailed in U.S. Provisional Patent Application Serial No. 60/193,767, filed on Mar. 31, 2000 and which is herein incorporated by reference.

In the present example, cancerous tumors derived from various human tissues were studied for PRO polypeptide-encoding gene expression relative to non-cancerous human tissue in an attempt to identify those PRO polypeptides which are overexpressed in cancerous tumors. Two sets of experimental data were generated. In one set, cancerous human colon tumor tissue and matched non-cancerous human colon tumor tissue from the same patient ("matched colon control") were obtained and analyzed for PRO polypeptide expression using the above described microarray technology. In the second set of data, cancerous human tumor tissue from any of a variety of different human tumors

was obtained and compared to a "universal" epithelial control sample which was prepared by pooling non-cancerous human tissues of epithelial origin, including liver, kidney, and lung. mRNA isolated from the pooled tissues represents a mixture of expressed gene products from these different tissues. Microarray hybridization experiments using the pooled control samples generated a linear plot in a 2-color analysis. The slope of the line generated in a 2-color analysis was then used to normalize the ratios of (test: control detection) within each experiment. The normalized ratios from various experiments were then compared and used to identify clustering of gene expression. Thus, the pooled "universal control" sample not only allowed effective relative gene expression determinations in a simple 2-sample comparison, it also allowed multi-sample comparisons across several experiments.

In the present experiments, nucleic acid probes derived from the herein described PRO polypeptide-encoding nucleic acid sequences were used in the creation of the microarray and RNA from the tumor tissues listed above were used for the hybridization thereto. A value based upon the normalized ratio:experimental ratio was designated as a "cutoff ratio". Only values that were above this cutoff ratio were determined to be significant. The results of these experiments are shown below, demonstrating that various PRO polypeptides of the present invention are significantly overexpressed in various human tumor tissues as compared to a non-cancerous human tissue control. As described above, these data demonstrate that the PRO polypeptides of the present invention are useful not only as diagnostic markers for the presence of one or more cancerous tumors, but also serve as therapeutic targets for the treatment of those tumors.

Molecule	is overexpressed in:	as compared to:
PRO3301	colon tumor	universal normal control
PRO3301	lung tumor	universal normal control
PRO3301	breast tumor	universal normal control
PRO3301	rectal tumor	universal normal control

Deposit of Material

The following materials have been deposited with the American Type Culture Collection, 10801 University Blvd., Manassas, Va. 20110-2209, USA (ATCC):

Material	ATCC Deposit No.	Deposit Date
DNA145887-2849	PTA-1532	March 21, 2000

These deposits were made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure and the Regulations thereunder (Budapest Treaty). This assures maintenance of a viable culture of the deposit for 30 years from the date of deposit. The deposits will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted availability of the progeny of the culture of the deposit to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S. or foreign patent application, whichever comes first, and assures availability of the progeny to one determined by the U.S. Commissioner of Patents and Trademarks to be entitled thereto according to

35 USC § 122 and the Commissioner's rules pursuant thereto (including 37 CFR § 1.14 with particular reference to 886 OG 638).

The assignee of the present application has agreed that if a culture of the materials on deposit should die or be lost or

destroyed when cultivated under suitable conditions, the materials will be promptly replaced on notification with another of the same. Availability of the deposited material is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its patent laws.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 7

<210> SEQ ID NO 1

<211> LENGTH: 1318

<212> TYPE: DNA

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 1

cagtttcttc atctgtaaca tcaaatgaat aataatacca atctcctaga	50
cttcataaga ggattaacaa agacaaaata tgggaaaaac ataacatggc	100
gtcccataat tattagatct tattattgac actaaaatgg cattaaaaatt	150
accaaaaagg agacagcatc tgtttcctct ttggtcctga gctgggttaa	200
aggaacactg gttgcctgaa cagtcacact tgcaaccatg atgcctaaac	250
attgctttct aggttcctc atcagtttct tccttactgg tgtagcagga	300
actcagtc aa cgcatgagtc tctgaagcct cagagggtag aatttcagtc	350
ccgaaatttt cacaacattt tgcaatggca gcctgggagg gcacttactg	400
gcaacagcag tgtctatttt gtgcagtaca aaatcatgtt ctcatgcagc	450
atgaaaagct ctcaccagaa gccaaagtga tgctggcagc acatttcttg	500
taacttccca ggctgcagaa cattggctaa atatggacag agacaatgga	550
aaaataaaga agactgttgg ggtactcaag aactctcttg tgaccttacc	600
agtgaacct cagacatata ggaaccttat tacgggaggg tgagggcggc	650
ctcggctggg agctactcag aatggagcat gacgccgcgg ttactccct	700
ggtgggaaac aaaaatagat cctccagtca tgaatataac ccaagtcaat	750
ggctctttgt tggtaatctt ccatgctcca aatttaccat atagatacca	800
aaaggaaaaa aatgtatcta tagaagatta ctatgaacta ctataccgag	850
ttttataat taacaattca ctagaaaagg agcaaaagg ttatgaagg	900
gtcacagag cggttgaat tgaagctcta acaccacact ccagctactg	950
tgtagtggt gaaatatatc agcccatgtt agacagaaga agtcagagaa	1000
gtgaagagag atgtgtggaa attccatgac ttgtggaatt tggcattcag	1050
caatgtggaa attctaaagc tccctgagaa caggatgact cgtgtttgaa	1100
ggatcttatt taaaattgtt ttgtatttt cttaaagcaa tattcactgt	1150
tacaccttgg ggacttcttt gtttatccat tcttttatcc tttatatttc	1200
atttgtaaac tatatttgaa cgacattccc cccgaaaaat tgaatgtaa	1250
agatgaggca gagaataaag tgttctatga aaaaaaaaaa aaaaaaaaaa	1300
aaaaaaaaa aaaaaaaaaa	1318

<210> SEQ ID NO 2

<211> LENGTH: 262

<212> TYPE: PRT

-continued

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 2

```

Met Pro Lys His Cys Phe Leu Gly Phe Leu Ile Ser Phe Phe Leu
 1           5           10          15
Thr Gly Val Ala Gly Thr Gln Ser Thr His Glu Ser Leu Lys Pro
          20          25          30
Gln Arg Val Gln Phe Gln Ser Arg Asn Phe His Asn Ile Leu Gln
          35          40          45
Trp Gln Pro Gly Arg Ala Leu Thr Gly Asn Ser Ser Val Tyr Phe
          50          55          60
Val Gln Tyr Lys Ile Met Phe Ser Cys Ser Met Lys Ser Ser His
          65          70          75
Gln Lys Pro Ser Gly Cys Trp Gln His Ile Ser Cys Asn Phe Pro
          80          85          90
Gly Cys Arg Thr Leu Ala Lys Tyr Gly Gln Arg Gln Trp Lys Asn
          95          100         105
Lys Glu Asp Cys Trp Gly Thr Gln Glu Leu Ser Cys Asp Leu Thr
          110         115         120
Ser Glu Thr Ser Asp Ile Gln Glu Pro Tyr Tyr Gly Arg Val Arg
          125         130         135
Ala Ala Ser Ala Gly Ser Tyr Ser Glu Trp Ser Met Thr Pro Arg
          140         145         150
Phe Thr Pro Trp Trp Glu Thr Lys Ile Asp Pro Pro Val Met Asn
          155         160         165
Ile Thr Gln Val Asn Gly Ser Leu Leu Val Ile Leu His Ala Pro
          170         175         180
Asn Leu Pro Tyr Arg Tyr Gln Lys Glu Lys Asn Val Ser Ile Glu
          185         190         195
Asp Tyr Tyr Glu Leu Leu Tyr Arg Val Phe Ile Ile Asn Asn Ser
          200         205         210
Leu Glu Lys Glu Gln Lys Val Tyr Glu Gly Ala His Arg Ala Val
          215         220         225
Glu Ile Glu Ala Leu Thr Pro His Ser Ser Tyr Cys Val Val Ala
          230         235         240
Glu Ile Tyr Gln Pro Met Leu Asp Arg Arg Ser Gln Arg Ser Glu
          245         250         255
Glu Arg Cys Val Glu Ile Pro
          260

```

<210> SEQ ID NO 3

<211> LENGTH: 27

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 3

ctggcaacag cagtgtctat tttgtgc

27

<210> SEQ ID NO 4

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 4

-continued

taagtgcctt cccaggctgc c	21
-------------------------	----

<210> SEQ ID NO 5
 <211> LENGTH: 52
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic oligonucleotide probe

<400> SEQUENCE: 5

tcctccagtc atgaatataa cccaagtcaa tggctctttg ttgtaattc	50
---	----

tc	52
----	----

<210> SEQ ID NO 6
 <211> LENGTH: 1705
 <212> TYPE: DNA
 <213> ORGANISM: Homo Sapien

<400> SEQUENCE: 6

tgaatgact tccacggctg ggacgggaac cttccaccca cagctatgcc	50
---	----

tctgattggt gaattggtgaa ggtgcctgtc taacttttct gtaaaaagaa	100
---	-----

ccagctgcct ccaggcagcc agccctcaag catcacttac aggaccagag	150
--	-----

ggacaagaca tgactgtgat gaggagctgc ttctgccaat ttaacaccaa	200
--	-----

gaagaattga ggctgcttgg gaggaaggcc aggaggaaca cgagactgag	250
--	-----

agatgaattt tcaacagagg ctgcaaagcc tgtggacttt agccagaccc	300
--	-----

ttctgccctc ctttgcctgg gacagcctct caaatgcaga tggttgtgct	350
--	-----

cccttgccctg ggttttaccg tgcttctctg gagccaggtg tcaggggccc	400
---	-----

agggccaaga attccacttt gggccctgcc aagtgaaggg ggttgttccc	450
--	-----

cagaaactgt gggaaagcctt ctgggctgtg aaagacacta tgcaagctca	500
---	-----

ggataaacatc acgagtgcct ggctgctgca gcaggaggtt ctgcagaacg	550
---	-----

tctcggatgc tgagagctgt taccttgtcc acaccctgct ggagttctac	600
--	-----

ttgaaaactg ttttcaaaaa ccaccacaat agaacagttg aagtcaggac	650
--	-----

tctgaagtca ttctctactc tggccaacaa ctttgttctc atcgtgtcac	700
--	-----

aactgcaacc cagtcaagaa aatgagatgt tttccatcag agacagtgca	750
--	-----

cacaggcggt ttctgctatt ccggagagca ttcaaacagt tggacgtaga	800
--	-----

agcagctctg accaaaagccc ttgggggaagt ggacattctt ctgacctgga	850
--	-----

tgcagaaatt ctacaagctc tgaatgtcta gaccaggacc tccctcccc	900
---	-----

tggcactggt ttgttccctg tgtcatttca aacagtctcc cttcctatgc	950
--	-----

tgttcaactg acacttcacg cccttggcca tgggtcccat tcttggccca	1000
--	------

ggattattgt caaagaagtc attctttaag cagcgccagt gacagtcagg	1050
--	------

gaagggtcct ctggatgctg tgaagagtct acagagaaga ttcttgtatt	1100
--	------

tattacaact ctatttaatt aatgtcagta tttcaactga agttctattt	1150
--	------

atttgtgaga ctgtaagtta catgaaggca gcagaatatt gtgccccatg	1200
--	------

cttctttacc cctcacaatc cttgccacag tgtggggcag tggatgggtg	1250
--	------

cttagtaagt acttaataaa ctgtgggtgct ttttttgcc tgcttttgga	1300
--	------

ttgttaaaaa acagagaggg atgcttggat gtaaaactga acttcagagc	1350
--	------

-continued

atgaaatca cactgtcttc tgatatctgc agggacagag cattggggtg	1400
ggggtaaggt gcatctgttt gaaaagtaaa cgataaaatg tggattaaag	1450
tgcccagcac aaagcagatc ctcaataaac atttcatttc ccacccacac	1500
tcgccagctc accccatcat cctttccct tggtgccctc cttttttttt	1550
tatcctagtc attcttcctt aatcttcac ttgagtgtca agctgacctt	1600
gctgatgggtg acattgcacc tggatgtact atccaatctg tgatgacatt	1650
ccctgctaataaaaagacaac ataactcaa aaaaaaaaaa aaaaaaaaaa	1700
aaaaa	1705

<210> SEQ ID NO 7

<211> LENGTH: 206

<212> TYPE: PRT

<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 7

Met Asn Phe Gln Gln Arg Leu Gln Ser Leu Trp Thr Leu Ala Arg	
1 5 10 15	
Pro Phe Cys Pro Pro Leu Leu Ala Thr Ala Ser Gln Met Gln Met	
20 25 30	
Val Val Leu Pro Cys Leu Gly Phe Thr Leu Leu Leu Trp Ser Gln	
35 40 45	
Val Ser Gly Ala Gln Gly Gln Glu Phe His Phe Gly Pro Cys Gln	
50 55 60	
Val Lys Gly Val Val Pro Gln Lys Leu Trp Glu Ala Phe Trp Ala	
65 70 75	
Val Lys Asp Thr Met Gln Ala Gln Asp Asn Ile Thr Ser Ala Arg	
80 85 90	
Leu Leu Gln Gln Glu Val Leu Gln Asn Val Ser Asp Ala Glu Ser	
95 100 105	
Cys Tyr Leu Val His Thr Leu Leu Glu Phe Tyr Leu Lys Thr Val	
110 115 120	
Phe Lys Asn His His Asn Arg Thr Val Glu Val Arg Thr Leu Lys	
125 130 135	
Ser Phe Ser Thr Leu Ala Asn Asn Phe Val Leu Ile Val Ser Gln	
140 145 150	
Leu Gln Pro Ser Gln Glu Asn Glu Met Phe Ser Ile Arg Asp Ser	
155 160 165	
Ala His Arg Arg Phe Leu Leu Phe Arg Arg Ala Phe Lys Gln Leu	
170 175 180	
Asp Val Glu Ala Ala Leu Thr Lys Ala Leu Gly Glu Val Asp Ile	
185 190 195	
Leu Leu Thr Trp Met Gln Lys Phe Tyr Lys Leu	
200 205	

What is claimed is:

1. An isolated nucleic acid molecule having at least 80% sequence identity to:

(a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;

(b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide;

(c) the nucleic acid sequence shown as SEQ ID NO:1;

(d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or

(e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532; wherein said isolated nucleic acid encodes a polypeptide which is a receptor for and binds to the ligand polypeptide shown as SEQ ID NO:4.

2. An isolated nucleic acid molecule having at least 85% sequence identity to:

- (a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;
 - (b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide;
 - (c) the nucleic acid sequence shown as SEQ ID NO:1;
 - (d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or
 - (e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532; wherein said isolated nucleic acid encodes a polypeptide which is a receptor for and binds to the ligand polypeptide shown as SEQ ID NO:4.
3. An isolated nucleic acid molecule having at least 90% sequence identity to:
- (a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;
 - (b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide;
 - (c) the nucleic acid sequence shown as SEQ ID NO:1;
 - (d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or
 - (e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532; wherein said isolated nucleic acid encodes a polypeptide which is a receptor for and binds to the ligand polypeptide shown as SEQ ID NO:4.
4. An isolated nucleic acid molecule having at least 95% sequence identity to:
- (a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;
 - (b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide;
 - (c) the nucleic acid sequence shown as SEQ ID NO:1;
 - (d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or
 - (e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532; wherein said isolated nucleic acid encodes a polypeptide which is a receptor for and binds to the ligand polypeptide shown as SEQ ID NO:4.
5. An isolated nucleic acid molecule having at least 99% sequence identity to:
- (a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;
 - (b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide;
 - (c) the nucleic acid sequence shown as SEQ ID NO:1;
 - (d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or

- (e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532; wherein said isolated nucleic acid encodes a polypeptide which is a receptor for and binds to the ligand polypeptide shown as SEQ ID NO:4.
6. An isolated nucleic acid molecule comprising:
- (a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;
 - (b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide;
 - (c) the nucleic acid sequence shown as SEQ ID NO:1;
 - (d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or
 - (e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532.
7. The isolated nucleic acid molecule of claim 6 comprising a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2.
8. The isolated nucleic acid molecule of claim 6 comprising a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide.
9. The isolated nucleic acid molecule of claim 6 comprising the nucleic acid sequence shown as SEQ ID NO:1.
10. The isolated nucleic acid molecule of claim 6 comprising the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1.
11. The isolated nucleic acid molecule of claim 6 comprising the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532.
12. An isolated nucleic acid molecule that hybridizes to:
- (a) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2;
 - (b) a nucleic acid sequence encoding the polypeptide shown as SEQ ID NO:2, lacking its associated signal peptide;
 - (c) the nucleic acid sequence shown as SEQ ID NO:1;
 - (d) the full-length coding sequence of the nucleic acid sequence shown as SEQ ID NO:1; or
 - (e) the full-length coding sequence of cDNA deposited under ATCC accession number PTA-1532; wherein said isolated nucleic acid encodes a polypeptide which is a receptor for and binds to the ligand polypeptide shown as SEQ ID NO:4.
13. A vector comprising the isolated nucleic acid molecule of claim 1.
14. The vector of claim 13, wherein said isolated nucleic acid is operably linked to control sequences recognized by a host cell transformed with the vector.
15. A host cell comprising the vector of claim 13.
16. The host cell of claim 15, wherein said cell is a CHO cell, an *E. coli* or a yeast cell.

* * * * *